



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### IL5 recombinant monoclonal antibody, clone 2E3

**Catalog Number:** RAB03587

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human IL5.

**Clone Name:** 2000

**Immunogen:** Original antibody is raised against recombinant protein corresponding to human IL5.

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Isotype:** IgG

**Recommend Usage:** ELISA

Flow Cytometry

Neutralization

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS (0.02% Proclin 300)

**Storage Instruction:** Store at 4°C for 3 months. For long term storage, aliquot and store at -20°C.

Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 3567

**Gene Symbol:** IL5

**Gene Alias:** EDF, IL-5, TRF

**Gene Summary:** The protein encoded by this gene is a cytokine that acts as a growth and differentiation factor for both B cells and eosinophils. This cytokine is a main regulator of eosinopoiesis, eosinophil maturation and activation. The elevated production of this cytokine is reported to be related to asthma or hypereosinophilic syndromes. The receptor of this cytokine is a

heterodimer, whose beta subunit is shared with the receptors for interleukine 3 (IL3) and colony stimulating factor 2 (CSF2/GM-CSF). This gene, together with those for interleukin 4 (IL4), interleukin 13 (IL13), and CSF2, form a cytokine gene cluster on chromosome 5. This cytokine, IL4, and IL13 are found to be regulated coordinately by long-range regulatory elements spread over 120 kilobases on chromosome 5q31. [provided by RefSeq]