



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

LTA recombinant monoclonal antibody, clone 3F12.2D3

Catalog Number: RAB03637

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human LTA.

Clone Name: 3F12.2D3

Immunogen: Original antibody is raised against purified recombinant human LTalpha expressed in *Escherichia coli*.

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Isotype: IgG

Recommend Usage: ELISA

Flow Cytometry

Neutralization

Surface Plasmon Resonance

Western Blot

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS (0.02% Proclin 300)

Storage Instruction: Store at 4°C for 3 months. For long term storage, aliquot and store at -20°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 4049

Gene Symbol: LTA

Gene Alias: LT, TNFB, TNFSF1

Gene Summary: The encoded protein, a member of the tumor necrosis factor family, is a cytokine produced by lymphocytes. The protein is highly inducible, secreted, and forms heterotrimers with lymphotoxin-beta which

anchor lymphotoxin-alpha to the cell surface. This protein also mediates a large variety of inflammatory, immunostimulatory, and antiviral responses, is involved in the formation of secondary lymphoid organs during development and plays a role in apoptosis. Genetic variations in this gene are associated with susceptibility to leprosy type 4 and psoriatic arthritis]