



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### **NFKB2 recombinant monoclonal antibody, clone SAIC-26C-15**

**Catalog Number:** RAB03718

**Regulatory Status:** For research use only (RUO)

**Product Description:** Mouse recombinant monoclonal antibody raised against synthetic peptide corresponding to human NFKB2.

**Clone Name:** SAIC-26C-15

**Immunogen:** Original antibody is raised against a synthetic peptide corresponding to human NFKB2

**Sequence:** IEVDLVTHSDPPR

**Antibody Species:** Mouse

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Conjugation:** Unconjugated

**Concentration:** batch dependent

**Isotype:** IgG1 kappa

**Recommend Usage:** Immuno-MRM

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS with 0.02% Proclin 300

**Storage Instruction:** Store at 4°C for up to 3 months.  
For longer storage, aliquot and store at -20°C.  
Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 4791

**Gene Symbol:** NFKB2

**Gene Alias:** LYT-10, LYT10

**Gene Summary:** NFKB has been detected in numerous cell types that express cytokines, chemokines, growth

factors, cell adhesion molecules, and some acute phase proteins in health and in various disease states. NFKB is activated by a wide variety of stimuli such as cytokines, oxidant-free radicals, inhaled particles, ultraviolet irradiation, and bacterial or viral products. Inappropriate activation of NF-kappa-B has been linked to inflammatory events associated with autoimmune arthritis, asthma, septic shock, lung fibrosis, glomerulonephritis, atherosclerosis, and AIDS. In contrast, complete and persistent inhibition of NF-kappa-B has been linked directly to apoptosis, inappropriate immune cell development, and delayed cell growth. For reviews, see Chen et al. (1999) [PubMed 9895331] and Baldwin (1996) [PubMed 8717528]. [supplied by OMIM]