

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

#### **Datasheet**

# NOTCH 1 recombinant monoclonal antibody, clone E6

Catalog Number: RAB03723

Regulatory Status: For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against a fusion protein consisting of EGF domains 1–12 of murine Notch1 fused to a human Fc domain and binds to to the NRR of mouse Notch1.

Clone Name: E6

**Immunogen:** Original antibody is raised against recombinant protein corresponding to a fusion protein consisting of EGF domains 1–12 of murine Notch1 fused to a human Fc domain and binds to to the NRR of mouse Notch1

Antibody Species: Rabbit

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Liquid

Conjugation: Unconjugated

Concentration: batch dependent

Isotype: IgG lambda

Recommend Usage: ELISA

Flow cytometry Immunofluorescence Immunohistochemistry

The optimal working dilution should be determined by

the end user.

Storage Buffer: In PBS with 0.02% Proclin 300

**Storage Instruction:** Store at 4°C for up to 3 months.

For longer storage, aliquot and store at -20°C. Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 4851

Gene Symbol: NOTCH1

Gene Alias: TAN1, hN1

Gene Summary: This gene encodes a member of the Notch family. Members of this Type 1 transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple, different domain types. Notch family members play a role in a variety of developmental processes by controlling cell fate decisions. The Notch signaling network is evolutionarily conserved intercellular signaling pathway which regulates interactions between physically adjacent cells. In Drosophilia, notch interaction with its cell-bound ligands (delta, serrate) establishes an intercellular signaling pathway that plays a key role in development. Homologues of the notch-ligands have also been identified in human, but precise interactions between these ligands and the human notch homologues remain to be determined. This protein is cleaved in the trans-Golgi network, and presented on the cell surface as a heterodimer. This protein functions as a receptor for membrane bound ligands, and may play multiple roles during development. [provided by RefSeq]