



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### PRDM4 recombinant monoclonal antibody, clone RAB-C367

**Catalog Number:** RAB03760

**Regulatory Status:** For research use only (RUO)

**Product Description:** Human recombinant monoclonal antibody raised against human PRDM4.

**Clone Name:** RAB-C367

**Immunogen:** Original antibody is raised against recombinant protein corresponding to human PRDM4

**Antibody Species:** Human

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Conjugation:** Unconjugated

**Concentration:** batch dependent

**Isotype:** Fab fragment kappa

**Recommend Usage:** ChIP

ELISA

Immunofluorescence

Immunoprecipitation-Mass Spectrometry

The optimal working dilution should be determined by the end user.

**Storage Buffer:** Lyophilized from PBS with 0.02% Proclin 300

**Storage Instruction:** Store at 4°C for 3 months. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 11108

**Gene Symbol:** PRDM4

**Gene Alias:** MGC45046, PFM1

**Gene Summary:** The protein encoded by this gene is a

transcription factor of the PR-domain protein family. It contains a PR-domain and multiple zinc finger motifs. Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. An elevated expression level of this gene has been observed in PC12 cells treated with nerve growth factor, beta polypeptide (NGF). This gene is located in a chromosomal region that is thought to contain tumor suppressor genes. [provided by RefSeq]