



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TNFRSF8 recombinant monoclonal antibody, clone Ki-1/1747R

Catalog Number: RAB03797

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human CD30 partial protein.

Clone Name: Ki-1/1747R

Immunogen: Original antibody is raised against recombinant protein corresponding to human CD30 partial protein

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Conjugation: Unconjugated

Purification: Protein A affinity chromatography

Concentration: 0.2 mg/mL

Isotype: IgG

Recommend Usage: Flow cytometry (0.5-1 ug/million cells in 0.1ml)

Immunofluorescence (1-2 ug/mL)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)(0.5-1 ug/mL for 30 min at RT)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, 0.1 mg/ml BSA, 0.05% sodium azide

Storage Instruction: Store at 2~8°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 943

Gene Symbol: TNFRSF8

Gene Alias: CD30, D1S166E, KI-1

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq]