



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CD79A recombinant monoclonal antibody, clone IGA/1688R

Catalog Number: RAB03804

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against synthetic peptide corresponding to amino acids 202-216 (GTYQDVGSLNIADVQ) of human CD79a.

Clone Name: IGA/1688R

Immunogen: Original antibody is raised against a synthetic peptide corresponding to amino acids 202-216 (GTYQDVGSLNIADVQ) of human CD79a

Sequence: GTYQDVGSLNIADVQ

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Conjugation: Unconjugated

Purification: Protein A affinity chromatography

Concentration: 0.2 mg/mL

Isotype: IgG

Recommend Usage: Flow cytometry (0.5-1 ug/million cells in 0.1ml)
Immunofluorescence (0.5-1 ug/mL)
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)(0.25-0.5 ug/mL for 30 min at RT)
The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, 0.1 mg/ml BSA, 0.05% sodium azide

Storage Instruction: Store at 2-8°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 973

Gene Symbol: CD79A

Gene Alias: IGA, MB-1

Gene Summary: The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component, surface immunoglobulin (Ig). Surface Ig non-covalently associates with two other proteins, Ig-alpha and Ig-beta, which are necessary for expression and function of the B-cell antigen receptor. This gene encodes the Ig-alpha protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq]