



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CHGA recombinant monoclonal antibody, clone CHGA/1773R

Catalog Number: RAB03809

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against recombinant human Chromogranin A protein.

Clone Name: CHGA/1773R

Immunogen: Original antibody is raised against recombinant protein corresponding to recombinant human Chromogranin A protein

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: Does not react with rat.

Form: Liquid

Conjugation: Unconjugated

Purification: Protein A affinity chromatography

Concentration: 0.2 mg/mL

Isotype: IgG

Recommend Usage: Flow cytometry (0.5-1 ug/million cells in 0.1mL)

Immunofluorescence (1-2 ug/mL)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)(0.25-0.5 ug/mL for 30 min at RT)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, 0.1 mg/ml BSA, 0.05% sodium azide

Storage Instruction: Store at 2~8°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 1113

Gene Symbol: CHGA

Gene Alias: CGA

Gene Summary: The protein encoded by this gene is a member of the chromogranin/secretogranin family of neuroendocrine secretory proteins. It is found in secretory vesicles of neurons and endocrine cells. This gene product is a precursor to three biologically active peptides; vasostatin, pancreastatin, and parastatin. These peptides act as autocrine or paracrine negative modulators of the neuroendocrine system. Other peptides, including chromostatin, beta-granin, WE-14 and GE-25, are also derived from the full-length protein. However, biological activities for these molecules have not been shown. [provided by RefSeq]