



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TLR2 recombinant monoclonal antibody, clone TL2.1

Catalog Number: RAB03908

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against immunising BALB/c mice with CHO cells transfected with human TLR2 cDNA.

Clone Name: TL2.1

Immunogen: Original antibody is raised against recombinant protein corresponding to immunising BALB/c mice with CHO cells transfected with human TLR2 cDNA.

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Isotype: IgG

Recommend Usage: Blocking

Flow cytometry

Immunofluorescence

Immunoprecipitation

Western Blot

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS (0.02% Proclin 300)

Storage Instruction: Store at 4°C for 3 months. For long term storage store at -20°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 7097

Gene Symbol: TLR2

Gene Alias: CD282, TIL4

Gene Summary: The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which

plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from *Drosophila* to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This gene is expressed most abundantly in peripheral blood leukocytes, and mediates host response to Gram-positive bacteria and yeast via stimulation of NF-kappaB. [provided by RefSeq]