



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

XPNPEP1 recombinant monoclonal antibody, clone SAIC-07E-1

Catalog Number: RAB03971

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against synthetic peptide corresponding to peptide "TSLSLDEVYLIDSGAQYK" derived from Xaa-Pro aminopeptidase 1 conjugated to KLH.

Clone Name: SAIC-07E-1

Immunogen: Original antibody is raised against recombinant protein corresponding to peptide "TSLSLDEVYLIDSGAQYK" derived from Xaa-Pro aminopeptidase 1 conjugated to KLH.

Sequence: TSLSLDEVYLIDSGAQYK

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Isotype: IgG kappa

Recommend Usage: ELISA

Flow cytometry

Immunofluorescence

Immuno-MRM

Western Blot

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS (0.02% Proclin 300)

Storage Instruction: Store at 4°C for 3 months. For long term storage store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 7511

Gene Symbol: XPNPEP1

Gene Alias: SAMP, XPNPEP, XPNPEPL, XPNPEPL1

Gene Summary: X-prolyl aminopeptidase (EC 3.4.11.9) is a proline-specific metalloaminopeptidase that specifically catalyzes the removal of any unsubstituted N-terminal amino acid that is adjacent to a penultimate proline residue. Because of its specificity toward proline, it has been suggested that X-prolyl aminopeptidase is important in the maturation and degradation of peptide hormones, neuropeptides, and tachykinins, as well as in the digestion of otherwise resistant dietary protein fragments, thereby complementing the pancreatic peptidases. Deficiency of X-prolyl aminopeptidase results in excretion of large amounts of imino-oligopeptides in urine (Blau et al., 1988 [PubMed 3141711]).[supplied by OMIM]