



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IKBKB recombinant monoclonal antibody, clone 5D12

Catalog Number: RAB03990

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human IKBKB.

Clone Name: 5D12

Immunogen: Original antibody is raised against recombinant protein corresponding to full length human IKBKB.

Theoretical MW (kDa): Calculated MW: 87, 8

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Affinity-chromatography

Isotype: IgG

Recommend Usage: ELISA

Western Blot(1:500-1:5000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH7.4 (150mM NaCl, 50% glycerol and 0.02% sodium azide)

Storage Instruction: store at -20 °C or -80 °C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3551

Gene Symbol: IKBKB

Gene Alias: FLJ40509, IKK-beta, IKK2, IKKB, MGC131801, NFKBIKB

Gene Summary: NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL (MIM 164910), RELA (MIM

164014), or RELB (MIM 604758) to form the NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, MIM 164008, or NFKBIB, MIM 604495), which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, MIM 600664, or IKBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).[supplied by OMIM]