



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### LOXL2 recombinant monoclonal antibody, clone 2E5

**Catalog Number:** RAB04127

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human, mouse and rat LOXL2.

**Clone Name:** 200000

**Immunogen:** Original antibody is raised against recombinant protein corresponding to full length human LOXL2.

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Affinity-chromatography

**Isotype:** IgG

**Recommend Usage:** ELISA

Flow Cytometry (1:20-1:200)

Immunohistochemistry (1:50-1:200)

Immunofluorescence(1:20-1:200)

Western Blot (1:500-1:5000)

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, pH7.4 (150 mM NaCl, 50% glycerol and 0.02% sodium azide)

**Storage Instruction:** Store at -20°C or -80°C. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 4017

**Gene Symbol:** LOXL2

**Gene Alias:** LOR2, WS9-14

**Gene Summary:** This gene encodes a member of the lysyl oxidase gene family. The prototypic member of the

family is essential to the biogenesis of connective tissue, encoding an extracellular copper-dependent amine oxidase that catalyses the first step in the formation of crosslinks in collagens and elastin. A highly conserved amino acid sequence at the C-terminus end appears to be sufficient for amine oxidase activity, suggesting that each family member may retain this function. The N-terminus is poorly conserved and may impart additional roles in developmental regulation, senescence, tumor suppression, cell growth control, and chemotaxis to each member of the family. [provided by RefSeq]