



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### Histone H4 (acetyl K5) recombinant monoclonal antibody, clone 2C9

**Catalog Number:** RAB04181

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human histone H4.

**Clone Name:** 2C9

**Immunogen:** Original antibody is raised against a synthetic acetylated peptide corresponding to residues surrounding K5 of human histone H4.

**Theoretical MW (kDa):** Calculated MW: 11 kD

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Affinity chromatography

**Isotype:** IgG

**Recommend Usage:** ELISA

Immunocytochemistry (1:50-1:500)

Immunofluorescence (1:30-1:200)

Western Blot (1:500-1:2000)

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, pH7.4 (150mM NaCl, 50% glycerol and 0.02% sodium azide)

**Storage Instruction:** store at -20 °C or -80 °C. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 8359

**Gene Symbol:** HIST1H4A

**Gene Alias:** H4/a, H4FA

**Gene Summary:** Histones are basic nuclear proteins

that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6. [provided by RefSeq]