



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### RAF1 (phospho S621) recombinant monoclonal antibody, clone 1C2

**Catalog Number:** RAB04273

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human RAF1.

**Clone Name:** 1C2

**Immunogen:** Original antibody is raised against a synthetic phosphopeptide corresponding to residues surrounding S621 of human RAF1.

**Theoretical MW (kDa):** Calculated MW: 73 kD

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Affinity chromatography

**Isotype:** IgG

**Recommend Usage:** ELISA

Immunofluorescence (1:20-1:200)

Western Blot (1:500-1:5000)

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, 150 mM NaCl, pH 7.4 (50% glycerol, 0.02% sodium azide)

**Storage Instruction:** Store at -20 °C or -80 °C. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 5894

**Gene Symbol:** RAF1

**Gene Alias:** CRAF, NS5, Raf-1, c-Raf

**Gene Summary:** This gene is the cellular homolog of viral raf gene (v-raf). The encoded protein is a MAP

kinase kinase kinase (MAP3K), which functions downstream of the Ras family of membrane associated GTPases to which it binds directly. Once activated, the cellular RAF1 protein can phosphorylate to activate the dual specificity protein kinases MEK1 and MEK2, which in turn phosphorylate to activate the serine/threonine specific protein kinases, ERK1 and ERK2. Activated ERKs are pleiotropic effectors of cell physiology and play an important role in the control of gene expression involved in the cell division cycle, apoptosis, cell differentiation and cell migration. Mutations in this gene are associated with Noonan syndrome 5 and LEOPARD syndrome 2. [provided by RefSeq]