



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### CDKN1B recombinant monoclonal antibody, clone 1G2

**Catalog Number:** RAB04334

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human CDKN1B.

**Clone Name:** 1G2

**Immunogen:** Original antibody is raised against a synthetic peptide corresponding to human CDKN1B.

**Theoretical MW (kDa):** Calculated MW: 23 kD

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Affinity chromatography

**Isotype:** IgG

**Recommend Usage:** ELISA

Immunofluorescence (1:20-1:200)

Immunohistochemistry (1:50-1:200)

Western Blot (1:500-1:5000)

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, pH7.4 (150mM NaCl, 50% glycerol and 0.02% sodium azide)

**Storage Instruction:** Store at -20 °C or -80 °C. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 1027

**Gene Symbol:** CDKN1B

**Gene Alias:** CDKN4, KIP1, MEN1B, MEN4, P27KIP1

**Gene Summary:** This gene encodes a cyclin-dependent kinase inhibitor, which shares a limited similarity with

CDK inhibitor CDKN1A/p21. The encoded protein binds to and prevents the activation of cyclin E-CDK2 or cyclin D-CDK4 complexes, and thus controls the cell cycle progression at G1. The degradation of this protein, which is triggered by its CDK dependent phosphorylation and subsequent ubiquitination by SCF complexes, is required for the cellular transition from quiescence to the proliferative state. [provided by RefSeq]