



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### CD19 recombinant monoclonal antibody, clone 3D3

order to decrease the threshold for antigen receptor-dependent stimulation. [provided by RefSeq]

**Catalog Number:** RAB04351

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human CD19.

**Clone Name:** 3D3

**Immunogen:** Original antibody is raised against a synthetic peptide corresponding to human CD19.

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Affinity chromatography

**Isotype:** IgG

**Recommend Usage:** ELISA

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, pH7.4 (150mM NaCl, 50% glycerol and 0.02% sodium azide)

**Storage Instruction:** Store at -20 °C or -80 °C. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 930

**Gene Symbol:** CD19

**Gene Alias:** B4, MGC12802

**Gene Summary:** Lymphocytes proliferate and differentiate in response to various concentrations of different antigens. The ability of the B cell to respond in a specific, yet sensitive manner to the various antigens is achieved with the use of low-affinity antigen receptors. This gene encodes a cell surface molecule which assembles with the antigen receptor of B lymphocytes in