



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

HSPA8 recombinant monoclonal antibody, clone 3G10

Catalog Number: RAB04358

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human HSPA8.

Clone Name: 3G10

Immunogen: Original antibody is raised against a synthetic peptide corresponding to human HSPA8.

Theoretical MW (kDa): Calculated MW: 71, 5

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Affinity chromatography

Isotype: IgG

Recommend Usage: ELISA
Western Blot (1:500-1:5000)
The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH7.4 (150mM NaCl, 50% glycerol and 0.02% sodium azide)

Storage Instruction: Store at -20 °C or -80 °C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3312

Gene Symbol: HSPA8

Gene Alias: HSC54, HSC70, HSC71, HSP71, HSP73, HSPA10, LAP1, MGC131511, MGC29929, NIP71

Gene Summary: The product encoded by this gene belongs to the heat shock protein 70 family which contains both heat-inducible and constitutively

expressed members. The latter are called heat-shock cognate proteins. This gene encodes a heat-shock cognate protein. This protein binds to nascent polypeptides to facilitate correct folding. It also functions as an ATPase in the disassembly of clathrin-coated vesicles during transport of membrane components through the cell. Two alternatively spliced variants have been characterized to date. [provided by RefSeq]