



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

BAG3 recombinant monoclonal antibody, clone 3D8

Catalog Number: RAB04377

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human BAG3.

Clone Name: 3D8

Immunogen: Original antibody is raised against a synthetic peptide corresponding to human BAG3.

Theoretical MW (kDa): Calculated MW: 62 kD

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Affinity chromatography

Isotype: IgG

Recommend Usage: ELISA
Immunohistochemistry (1:50-1:200)
Western Blot (1:500-1:5000)
The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH7.4 (150 mM NaCl, 50% glycerol and 0.02% sodium azide)

Storage Instruction: Store at -20°C or -80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 9531

Gene Symbol: BAG3

Gene Alias: BAG-3, BIS, CAIR-1, MGC104307

Gene Summary: BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an

approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The protein encoded by this gene contains a WW domain in the N-terminal region and a BAG domain in the C-terminal region. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. [provided by RefSeq]