

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

PRKCQ recombinant monoclonal antibody, clone R05-5I4

Catalog Number: RAB04470

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal

antibody raised against human PRKCQ.

Clone Name: R05-5I4

Immunogen: Original antibody is raised against a synthetic peptide corresponding to human PRKCQ.

Theoretical MW (kDa): Calculated MW: 82 kD

Antibody Species: Rabbit

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Liquid

Purification: Affinity chromatography

Isotype: IgG

Recommend Usage: Western Blot (1:500-1:1000) The optimal working dilution should be determined by

the end user.

Storage Buffer: In 50mM Tris-Glycine, 150mM NaCl, pH 7.4 (40% glycerol, 0.05% BSA and 0.01% Sodium

azide)

Storage Instruction: Store at 4°C. For long term

storage store at -20°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 5588

Gene Symbol: PRKCQ

Gene Alias: MGC126514, MGC141919, PRKCT, nPKC-

theta

Gene Summary: Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can

be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role. The protein encoded by this gene is one of the PKC family members. It is a calcium-independent and phospholipid-dependent protein kinase. This kinase is important for T-cell activation. It is required for the activation of the transcription factors NF-kappaB and AP-1, and may link the T cell receptor (TCR) signaling complex to the activation of the transcription factors. [provided by RefSeq1