

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



#### www.abnova.com

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## Datasheet

# FGFR3 recombinant monoclonal antibody, clone R03-4E4

Catalog Number: RAB06522

Regulatory Status: For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human FGFR3.

Clone Name: R03-4E4

**Immunogen:** Original antibody is raised against recombinant protein corresponding to human FGFR3.

Theoretical MW (kDa): Calculated MW: 88 kD

Antibody Species: Rabbit

**Protocols:** See our web site at http://www.abnova.com/support/protocols.asp or product page for detailed protocols

Form: Liquid

Purification: Affinity purification

Isotype: IgG

**Recommend Usage:** Western Blot (1:500-1:1000) The optimal working dilution should be determined by the end use.

**Storage Buffer:** In PBS, 150 mM NaCl, pH 7.4 (50% glycerol and 0.02% Sodium azide)

**Storage Instruction:** Store at 4°C. For long term storage store at -20°C. Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 2261

Gene Symbol: FGFR3

Gene Alias: ACH, CD333, CEK2, HSFGFR3EX, JTK4

**Gene Summary:** This gene encodes a member of the fibroblast growth factor receptor (FGFR) family, with its amino acid sequence being highly conserved between members and among divergent species. FGFR family

members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream ultimately influencing mitogenesis signals, and differentiation. This particular family member binds acidic and basic fibroblast growth hormone and plays a role in bone development and maintenance. Mutations in this gene lead to craniosynostosis and multiple types of skeletal dysplasia. Three alternatively spliced transcript variants that encode different protein isoforms have been described. [provided by RefSeq]