



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

HDAC2 recombinant monoclonal antibody, clone 9H4

Catalog Number: RAB07405

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human HDAC2.

Clone Name: 9H4

Immunogen: Original antibody is raised against a synthetic peptide corresponding to human HDAC2.

Theoretical MW (kDa): Calculated MW: 56, 5

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Affinity chromatography purification

Isotype: IgG

Recommend Usage: ELISA

Immunofluorescence (1:20-1:200)

Immunohistochemistry (1:50-1:200)

Western Blot (1:500-1:5000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH7.4 (150 mM NaCl, 0.02% sodium azide and 50% glycerol)

Storage Instruction: Store at -20°C or -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3066

Gene Symbol: HDAC2

Gene Alias: RPD3, YAF1

Gene Summary: This gene product belongs to the histone deacetylase family. Histone deacetylases act via

the formation of large multiprotein complexes and are responsible for the deacetylation of lysine residues on the N-terminal region of the core histones (H2A, H2B, H3 and H4). This protein also forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus it plays an important role in transcriptional regulation, cell cycle progression and developmental events. [provided by RefSeq]