



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

NONO recombinant monoclonal antibody, clone 8C11

Catalog Number: RAB07472

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human NONO.

Clone Name: 8C11

Immunogen: Original antibody is raised against a synthetic peptide corresponding to human NONO.

Theoretical MW (kDa): Calculated MW: 55, 4

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Affinity chromatography purification

Isotype: IgG

Recommend Usage: ELISA

Immunofluorescence (1:20-1:200)

Immunohistochemistry (1:50-1:200)

Western Blot (1:500-1:5000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH7.4 (150 mM NaCl, 0.02% sodium azide and 50% glycerol)

Storage Instruction: Store at -20°C or -80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 4841

Gene Symbol: NONO

Gene Alias: NMT55, NRB54, P54, P54NRB

Gene Summary: This gene encodes an RNA-binding protein which plays various roles in the nucleus,

including transcriptional regulation and RNA splicing. A rearrangement between this gene and the transcription factor E3 gene has been observed in papillary renal cell carcinoma. Alternatively spliced transcript variants have been described. Pseudogenes exist on Chromosomes 2 and 16. [provided by RefSeq]