



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### PTGS2 recombinant monoclonal antibody, clone 10B2

**Catalog Number:** RAB07487

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human PTGS2.

**Clone Name:** 10B2

**Immunogen:** Original antibody is raised against a synthetic peptide corresponding to human PTGS2.

**Theoretical MW (kDa):** 69

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Affinity chromatography purification

**Isotype:** IgG

**Recommend Usage:** ELISA

Immunofluorescence (1:20-1:200)

Western Blot (1:500-1:5000)

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS, pH7.4 (150 mM NaCl, 0.02% sodium azide and 50% glycerol)

**Storage Instruction:** Store at -20°C or -80°C.  
Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 5743

**Gene Symbol:** PTGS2

**Gene Alias:** COX-2, COX2, GRIPGHS, PGG/HS, PGHS-2, PHS-2, hCox-2

**Gene Summary:** Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key

enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis. [provided by RefSeq]