



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

DMEM/F12 Medium

Storage Instruction: Store at 2-8°C. Protect from light.

Catalog Number: U0606

Regulation Status: For research use only (RUO)

Product Description: DMEM/F12 Medium is used in the culture of many mammalian cells, the basis for the development of serum-free medium, and is suitable for mammalian cell culture and clone density culture under low serum content.

Formulation: Contains 2.5 mM L-glutamine, 3151 mg/L D-glucose, 0.5 mM sodium pyruvate, and 8.1 mg/L phenol red.
No HEPES.
No sodium bicarbonate.

Medium Preparation: 1. The net weight of this product is 12.04 g, which can prepare 1 L DMEM/F12 culture medium.
2. Prepare about 900 mL ultrapure water in a container, then add a bag of DMEM/F12 Medium into the container, and wash out the residual powder in the bag with 50 mL ultrapure water.
3. Stir for 30 mins to dissolve all ingredients completely. After the solution is clear, add 2.483 g NaHCO₃ and continue stirring for 5-10 mins until dissolved.
4. Add ultrapure water to adjust the volume to 1 L.
5. If necessary, adjust the pH to 7.20-7.30 with 1 mol/L NaOH solution or 1 mol/L HCl solution. Since filtration will slightly increase the pH, the pH value here is lower than the target pH value (7.20-7.40).
6. The prepared solution should be sterilized using a 0.2 µm pore size filter membrane under positive pressure (ensure aseptic technique).
7. After filtration, a small amount of liquid culture medium can be taken for quality inspection, only after it passes the test can it be used further. When stored at 2-8°C with shading light, the liquid medium has a shelf life of 1 year.

Regulatory Status: For research use only (RUO)

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized