



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0


F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Product Number	ARP54491_P050-FITC
Product Page	<a href="http://www.avivasysbio.com/narf-antibody-middle-region-fitc-arp54491-p050-fitc.html">www.avivasysbio.com/narf-antibody-middle-region-fitc-arp54491-p050-fitc.html</a>
Name	NARF Antibody - middle region : FITC (ARP54491_P050-FITC)
Protein Size (# AA)	397 amino acids
Molecular Weight	45kDa
Conjugation	FITC: Fluorescein Isothiocyanate
NCBI Gene Id	26502
Host	Rabbit
Clonality	Polyclonal
Concentration	0.5 mg/ml
Gene Full Name	Nuclear prelamin A recognition factor
Alias Symbols	IOP2
Peptide Sequence	Synthetic peptide located within the following region: <a href="#">RGQAQTPDGHADKALLRQMEGIYADIPVRRPESSAHVQELYQEWLEGINS</a>
Product Format	Liquid. Purified antibody supplied in 1x PBS buffer.
Description of Target	Several proteins have been found to be prenylated and methylated at their carboxyl-terminal ends. Prenylation was initially believed to be important only for membrane attachment. However, another role for prenylation appears to be its importance in protein-protein interactions. The only nuclear proteins known to be prenylated in mammalian cells are prelamin A- and B-type lamins. Prelamin A is farnesylated and carboxymethylated on the cysteine residue of a carboxyl-terminal CaaX motif. This post-translationally modified cysteine residue is removed from prelamin A when it is endoproteolytically processed into mature lamin A. NARF binds to the prenylated prelamin A carboxyl-terminal tail domain. It may be a component of a prelamin A endoprotease complex. NARF is located in the nucleus, where it partially colocalizes with the nuclear lamina. It shares limited sequence similarity with iron-only bacterial hydrogenases.
Protein Interactions	UBC; LEF1; STAT5A; SHFM1; MMS19; CBX5; APP; LMNA;
Reconstitution and Storage	All conjugated antibodies should be stored in light-protected vials or covered with a light protecting material (i.e. aluminum foil). Conjugated antibodies are stable for at least 12 months at 4C. If longer storage is desired (24 months), conjugates may be diluted with up to 50% glycerol and stored at -20C to -80C. Freezing and thawing conjugated antibodies will compromise enzyme activity as well as antibody binding.
Datasheets/Manuals	Printable datasheet for <a href="#">anti-NARF (ARP54491_P050-FITC) antibody</a>
Blocking Peptide	For anti-NARF (ARP54491_P050-FITC) antibody is <a href="#">Catalog # AAP54491</a> (Previous Catalog # AAPP44502)
Immunogen	The immunogen is a synthetic peptide directed towards the middle region of human NARF
Uniprot ID	<a href="#">B3KPX2</a>
Protein Name	cDNA FLJ32404 fis, clone SKMUS2000380, highly similar to Homo sapiens nuclear prelamin A recognition factor (NARF), transcript variant 1, mRNA EMBL BAG51834.1
Protein Accession #	<a href="#">NP_001033707</a>
Purification	Affinity Purified
Nucleotide Accession #	<a href="#">NM_001038618</a>
Gene Symbol	<a href="#">NARF</a>
Predicted Species Reactivity	Human, Mouse, Rat, Cow, Dog, Guinea Pig, Horse, Rabbit
Application	WB

<b>Predicted Homology Based on Immunogen Sequence</b>	Cow: 85%; Dog: 100%; Guinea Pig: 92%; Horse: 85%; Human: 100%; Mouse: 92%; Rabbit: 85%; Rat: 100%
<b>Image 1</b>	

AVIVA SYSTEMS BIOLOGY manufactures and sells quality antibody products covering genome wide proteins.

This product is for Research Use Only. Not for diagnostic, human, or veterinary use.

Optimal conditions of its use should be determined by end users.

---

AVIVA SYSTEMS BIOLOGY

6370 Nancy Ridge Dr., Suite 104, San Diego, CA 92121 USA | Tel: (858)552-6979 | [info@avivasysbio.com](mailto:info@avivasysbio.com)