

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





ARG2 Antibody - C-terminal region : FITC (ARP54567_P050-FITC)

Data Sheet

Product Number	ARP54567_P050-FITC
Product Page	www.avivasysbio.com/arg2-antibody-c-terminal-region-fitc-arp54567-p050-fitc.html
Name	ARG2 Antibody - C-terminal region : FITC (ARP54567_P050-FITC)
Protein Size (# AA)	354 amino acids
Molecular Weight	39kDa
Conjugation	FITC: Fluorescein Isothiocyanate
NCBI Gene Id	384
Host	Rabbit
Clonality	Polyclonal
Concentration	0.5 mg/ml
Gene Full Name	Arginase, type II
Alias Symbols	-
Peptide Sequence	Synthetic peptide located within the following region: SALDLVEVNPQLATSEEAKTTANLAVDVIASSFGQTREGGHIVYDQLPT
Product Format	Liquid. Purified antibody supplied in 1x PBS buffer.
Reference	Mumenthaler, S.M., (2008) Int. J. Oncol. 32 (2), 357-365
Description of Target	Arginase catalyzes the hydrolysis of arginine to ornithine and urea. At least two isoforms of mammalian arginase exists (types I and II) which differ in their tissue distribution, subcellular localization, immunologic crossreactivity and physiologic function. ARG2 (type II isoform) is located in the mitochondria and expressed in extra-hepatic tissues, especially kidney. The physiologic role of this isoform is poorly understood; it is thought to play a role in nitric oxide and polyamine metabolism. Arginase catalyzes the hydrolysis of arginine to ornithine and urea. At least two isoforms of mammalian arginase exists (types I and II) which differ in their tissue distribution, subcellular localization, immunologic crossreactivity and physiologic function. The type II isoform encoded by this gene, is located in the mitochondria and expressed in extra-hepatic tissues, especially kidney. The physiologic role of this isoform is poorly understood; it is thought to play a role in nitric oxide and polyamine metabolism. Transcript variants of the type II gene resulting from the use of alternative polyadenylation sites have been described. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.
Protein Interactions	UBC; ARG1;
Reconstitution and Storage	All conjugated antibodies should be stored in light-protected vials or covered with a light protecting material (i.e. aluminum foil). Conjugated antibodies are stable for at least 12 months at 4C. If longer storage is desired (24 months), conjugates may be diluted with up to 50% glycerol and stored at -20C to -80C. Freezing and thawing conjugated antibodies will compromise enzyme activity as well as antibody binding.
Datasheets/Manuals	Printable datasheet for anti-ARG2 (ARP54567_P050-FITC) antibody
Blocking Peptide	For anti-ARG2 (ARP54567_P050-FITC) antibody is Catalog # AAP54567 (Previous Catalog # AAPP31351)
Immunogen	The immunogen is a synthetic peptide directed towards the C terminal region of human ARG2
Uniprot ID	<u>P78540</u>
Protein Name	Arginase-2, mitochondrial
Publications	Lukasova, M., Malaval, C., Gille, A., Kero, J. & Offermanns, S. Nicotinic acid inhibits progression of atherosclerosis in mice through its receptor GPR109A expressed by immune cells. J. Clin. Invest. 121, 1163-73 (2011). WB, IHC, Dog, Bovine, Pig, Rat, Mouse, Human, Rabbit, Guinea pig, Horse 21317532
Sample Type Confirmation	ARG2 is supported by BioGPS gene expression data to be expressed in Jurkat

Protein Accession #	<u>NP_001163</u>
Purification	Affinity Purified
Nucleotide Accession#	<u>NM_001172</u>
Gene Symbol	ARG2
Predicted Species Reactivity	Human, Mouse, Rat, Cow, Dog, Guinea Pig, Horse, Pig, Rabbit
Application	WB
Predicted Homology Based on Immunogen Sequence	Cow: 100%; Dog: 100%; Guinea Pig: 93%; Horse: 87%; Human: 100%; Mouse: 100%; Pig: 100%; Rabbit: 93%; Rat: 100%
Image 1	

AVIVA SYSTEMS BIOLOGY manufactures and sells quality antibody products covering genome wide proteins.

This product is for Research Use Only. Not for diagnostic, human, or veterinary use. Optimal conditions of its use should be determined by end users.

AVIVA SYSTEMS BIOLOGY

6370 Nancy Ridge Dr., Suite 104, San Diego, CA 92121 USA | Tel: (858)552-6979 | info@avivasysbio.com