

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



PSME2 Antibody - middle region : FITC (ARP56485_P050-FITC)

Data Sheet

ARP56485_P050-FITC
www.avivasysbio.com/psme2-antibody-middle-region-fitc-arp56485-p050-fitc.html
PSME2 Antibody - middle region : FITC (ARP56485_P050-FITC)
239 amino acids
27kDa
2
FITC: Fluorescein Isothiocyanate
5721
Rabbit
Polyclonal
0.5 mg/ml
Proteasome (prosome, macropain) activator subunit 2 (PA28 beta)
PA28B, REGbeta, PA28beta
Synthetic peptide located within the following region: QEKVLERVNAVKTKVEAFQTTISKYFSERGDAVAKASKETHVMDYRALVH
Liquid. Purified antibody supplied in 1x PBS buffer.
The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. The immunoproteasome contains an alternate regulator, referred to as the 11S regulator or PA28, that replaces the 19S regulator. Three subunits (alpha, beta and gamma) of the 11S regulator have been identified. This gene encodes the beta subunit of the 11S regulator, one of the two 11S subunits that is induced by gamma-interferon. Three beta and three alpha subunits combine to form a heterohexameric ring. Six pseudogenes have been identified on chromosomes 4, 5, 8, 10 and 13.
NAA10; PSMD14; SHFM1; UBC; PSME2; PSME1; PARK2; BAG3; NHLH1; MYOD1; EPHA2; CFTR; DDTL; TNN; CAP1; PTMA; PSMD7; PSMD2; PSMD1; PSMC5; PSMC4; PSMC2; PSMB5; PSMB3; PSMB1; PSMA6; PSMA5; PSMA3; PSMA2; PSMA1; GARS; APP; PSMC1; PSMD6; PSMB10; PSMB9; PSMB8; PSMB7;
All conjugated antibodies should be stored in light-protected vials or covered with a light protecting material (i.e. aluminum foil). Conjugated antibodies are stable for at least 12 months at 4C. If longer storage is desired (24 months), conjugates may be diluted with up to 50% glycerol and stored at -20C to -80C. Freezing and thawing conjugated antibodies will compromise enzyme activity as well as antibody binding.
Printable datasheet for anti-PSME2 (ARP56485_P050-FITC) antibody
For anti-PSME2 (ARP56485_P050-FITC) antibody is Catalog # AAP56485 (Previous Catalog # AAPP38940)
The immunogen is a synthetic peptide directed towards the middle region of human PSME2
Q9UL46
Proteasome activator complex subunit 2
<u>NP_002809</u>
1
Affinity Purified
Affinity Purified NM 002818

Predicted Species Reactivity	Human, Mouse, Rat, Cow, Dog, Guinea Pig, Horse, Pig, Rabbit, Zebrafish
Application	WB
Predicted Homology Based on Immunogen Sequence	Cow: 100%; Dog: 100%; Guinea Pig: 93%; Horse: 93%; Human: 100%; Mouse: 100%; Pig: 100%; Rabbit: 93%; Rat: 100%; Zebrafish: 79%
Image 1	

AVIVA SYSTEMS BIOLOGY manufactures and sells quality antibody products covering genome wide proteins.

This product is for Research Use Only. Not for diagnostic, human, or veterinary use. Optimal conditions of its use should be determined by end users.

AVIVA SYSTEMS BIOLOGY 6370 Nancy Ridge Dr., Suite 104, San Diego, CA 92121 USA | Tel: (858)552-6979 | info@avivasysbio.com