



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0


F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Product Number	ARP58175_P050-Biotin
Product Page	<a href="http://www.avivasysbio.com/psmc3-antibody-n-terminal-region-biotin-arp58175-p050-biotin.html">www.avivasysbio.com/psmc3-antibody-n-terminal-region-biotin-arp58175-p050-biotin.html</a>
Name	PSMC3 Antibody - N-terminal region : Biotin (ARP58175_P050-Biotin)
Protein Size (# AA)	439 amino acids
Molecular Weight	49kDa
Subunit	6A
Conjugation	Biotin
NCBI Gene Id	5702
Host	Rabbit
Clonality	Polyclonal
Concentration	0.5 mg/ml
Gene Full Name	Proteasome (prosome, macropain) 26S subunit, ATPase, 3
Alias Symbols	RPT5, TBP1
Peptide Sequence	Synthetic peptide located within the following region: <a href="#">IQIWVRQQHPESGEEAVALVEDLQKEPGRQRLEPCLMWLWEFLQRRAGVA</a>
Product Format	Liquid. Purified antibody supplied in 1x PBS buffer.
Reference	Pollice,A., (2007) Oncogene 26 (35), 5154-5162
Description of Target	<p>PSMC3 is a subunit of the 26S proteasome. 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. This subunit may compete with PSMC2 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex. The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases that have chaperone-like activity. This subunit may compete with PSMC2 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex. A pseudogene has been identified on chromosome 9. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.</p>
Protein Interactions	AMOTL2; UBC; PSMD9; PSMC6; PSMC3; KDM1A; HUWE1; SUMO2; PSMD14; MDM2; ASB11; UCHL5; PSMD3; PSMD2; PSMD1; PSMC2; PSMC1; ADRM1; JKAMP; FBXO6; PARK2; BAG3; GADD45A; NOS2; MYC; PSMD13; PSMC4; PSMA8; UBF1; ECD; STIP1; RANBP9; PSMD6; ZFYVE16; USP14; RUVBL1; PSM
Reconstitution and Storage	All conjugated antibodies should be stored in light-protected vials or covered with a light protecting material (i.e. aluminum foil). Conjugated antibodies are stable for at least 12 months at 4C. If longer storage is desired (24 months), conjugates may be diluted with up to 50% glycerol and stored at -20C to -80C. Freezing and thawing conjugated antibodies will compromise enzyme activity as well as antibody binding.
Datasheets/Manuals	Printable datasheet for <a href="#">anti-PSMC3 (ARP58175_P050-Biotin) antibody</a>

<b>Blocking Peptide</b>	For anti-PSMC3 (ARP58175_P050-Biotin) antibody is <a href="#">Catalog # AAP58175</a> (Previous Catalog # AAPP32627)
<b>Immunogen</b>	The immunogen is a synthetic peptide directed towards the N terminal region of human PSMC3
<b>Uniprot ID</b>	<a href="#">P17980</a>
<b>Protein Name</b>	26S protease regulatory subunit 6A
<b>Protein Accession #</b>	<a href="#">NP_002795</a>
<b>Purification</b>	Affinity Purified
<b>Nucleotide Accession #</b>	<a href="#">NM_002804</a>
<b>Gene Symbol</b>	<a href="#">PSMC3</a>
<b>Predicted Species Reactivity</b>	Human, Mouse, Rat, Cow, Dog, Guinea Pig, Horse, Rabbit, Zebrafish
<b>Application</b>	WB
<b>Predicted Homology Based on Immunogen Sequence</b>	Cow: 100%; Dog: 100%; Guinea Pig: 100%; Horse: 100%; Human: 100%; Mouse: 100%; Rabbit: 100%; Rat: 100%; Zebrafish: 100%
<b>Image 1</b>	

AVIVA SYSTEMS BIOLOGY manufactures and sells quality antibody products covering genome wide proteins.

This product is for Research Use Only. Not for diagnostic, human, or veterinary use.

Optimal conditions of its use should be determined by end users.

---

AVIVA SYSTEMS BIOLOGY

6370 Nancy Ridge Dr., Suite 104, San Diego, CA 92121 USA | Tel: (858)552-6979 | [info@avivasysbio.com](mailto:info@avivasysbio.com)