

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



## H2AFY Antibody - N-terminal region : FITC (ARP58284\_P050-FITC)

Data Sheet

Product Number	ARP58284_P050-FITC
Product Page	www.avivasysbio.com/h2afy-antibody-n-terminal-region-fitc-arp58284-p050-fitc.html
Name	H2AFY Antibody - N-terminal region : FITC (ARP58284_P050-FITC)
Protein Size (# AA)	371 amino acids
Molecular Weight	39kDa
Conjugation	FITC: Fluorescein Isothiocyanate
NCBI Gene Id	9555
Host	Rabbit
Clonality	Polyclonal
Concentration	0.5 mg/ml
Gene Full Name	H2A histone family, member Y
Alias Symbols	H2A.y, H2A/y, H2AFY, mH2A1, H2AF12M, MACROH2A1.1, macroH2A1.2
Peptide Sequence	Synthetic peptide located within the following region: HPKYRIGVGAPVYMAAVLEYLTAEILELAGNAARDNKKGRVTPRHILLAV
Product Format	Liquid. Purified antibody supplied in 1x PBS buffer.
Reference	Bernstein, E., (2008) Proc. Natl. Acad. Sci. U.S.A. 105 (5), 1533-1538
Description of Target	<ul> <li>Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. H2AFY is a member of the histone H2A family. It replaces conventional H2A histones in a subset of nucleosomes where it represses transcription and participates in stable X chromosome inactivation. Alternative splicing results in multiple transcript variants encoding different isoforms. Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a member of the histone H2A family. It replaces conventional H2A histones in a subset of nucleosomes where it represses transcription and participates in stable X chromosome inactivation. Alternative splicing results in multiple transcript variants encoding different isoforms.</li> <li>ERICH2; UBC; SUMO2; SUMO3; HIST1H3A; H3F3A; SUZ12; EED; RNF2; EZH2; BMI1; SRPK1; LMNA:</li> </ul>
Protein Interactions	ATF2; FN1; HIST1H3B; CENPA; APP; CBX5; CBX3; DIDO1; UBD; ATXN1L; NEDD8; ERBB2; SUMO1; HDGF; HDAC2; HDAC1; PARP1; tat; CUL3; SPOP; WDR77; PHF14; DEK; XRCC5; TOP1; RAN; H
Reconstitution and Storage	All conjugated antibodies should be stored in light-protected vials or covered with a light protecting material (i.e. aluminum foil). Conjugated antibodies are stable for at least 12 months at 4C. If longer storage is desired (24 months), conjugates may be diluted with up to 50% glycerol and stored at -20C to -80C. Freezing and thawing conjugated antibodies will compromise enzyme activity as well as antibody binding.
Datasheets/Manuals	Printable datasheet for anti-H2AFY (ARP58284_P050-FITC) antibody
Blocking Peptide	For anti-H2AFY (ARP58284_P050-FITC) antibody is <u>Catalog # AAP58284</u> (Previous Catalog # AAPP32883)
Immunogen	The immunogen is a synthetic peptide directed towards the N terminal region of human H2AFY
Uniprot ID	<u>075367</u>
Protein Name	Core histone macro-H2A.1
Sample Type Confirmation	H2AFY is strongly supported by BioGPS gene expression data to be expressed in MCF7

Protein Accession #	<u>NP_001035248</u>
Purification	Affinity Purified
Nucleotide Accession #	<u>NM_001040158</u>
Gene Symbol	H2AFY
Predicted Species Reactivity	Human, Mouse, Rat, Cow, Dog, Guinea Pig, Horse, Rabbit, Yeast, Zebrafish
Application	WB
Predicted Homology Based on Immunogen Sequence	Cow: 100%; Dog: 100%; Guinea Pig: 100%; Horse: 100%; Human: 100%; Mouse: 100%; Rabbit: 100%; Rat: 100%; Yeast: 93%; Zebrafish: 100%
Image 1	<b>N</b>

AVIVA SYSTEMS BIOLOGY manufactures and sells quality antibody products covering genome wide proteins.

This product is for Research Use Only. Not for diagnostic, human, or veterinary use. Optimal conditions of its use should be determined by end users.

AVIVA SYSTEMS BIOLOGY 6370 Nancy Ridge Dr., Suite 104, San Diego, CA 92121 USA | Tel: (858)552-6979 | info@avivasysbio.com