

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



## Goat Anti-Rat IgG(H+L) Antibody (OASB01922)

Data Sheet

Product Number	OASB01922
Product Page	www.avivasysbio.com/rat-igg-h-l-mouse-ads-antibody-oasb01922.html
Name	Goat Anti-Rat IgG(H+L) Antibody (OASB01922)
Isotype	IgG
Host	Goat
Clonality	Polyclonal
Concentration	1.0 mg/mL
Gene Full Name	Ig gamma-1 chain C region
Reconstitution and Storage	Store at 2-8C
Storage	<ul> <li>The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of 100 mM boratebuffered saline, pH 8.2. No preservatives or amine- containing buffer salts added. Store at 2- 8 C</li> <li>The fluorescein (FITC), rhodamine (TRITC), and Texas Red (TXRD) conjugates are supplied as 1.0 mg in1.0 mL PBS/NaN3. Store at 2- 8 C</li> <li>The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL of stock solution in 50mM Tris/1mMMgCl2/50% Glycerol, pH 8.0, containing 0.1% NaN3 as preservative. Store at 2- 8 C or long- term at - 20 C</li> <li>The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50%PBS, pH 7.4. No preservative added. Store at 2- 8 C or long- term at - 20 C</li> <li>The Beta- galactosidase (BGAL) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50%PBS, glycerol/50%PBS/NaN3 as preservative. Store at 2- 8 C or long- term at - 20 C</li> <li>The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL PBS/NaN3. Store at 2- 8 C</li> <li>The R- phycoerythrin (R- PE) conjugate is supplied as 0.5 mg in 1.0 mL or 0.25 mg in 0.5 mL of PBS/NaN3 as a stabilizing agent. Store at 2- 8 C. Do not freeze!</li> <li>Protect fluorochrome- conjugated forms from light. Reagents are stable for the period shown on the label ifstored as directed.</li> </ul>
Other Applications Data	Since applications vary, you should determine the optimum workingdilution of the product that is appropriate for your specific need.
Datasheets/Manuals	Printable datasheet for anti-IGHG1 (OASB01922) antibody
Specificity	IgG(H+L)
Additional Information	<b>Description:</b> Pooled antisera from goats hyperimmunized with rat IgG; purified by affinity chromatography on pooled rat IgG covalently linked to agarose; reacts with the heavy and light chains of rat IgG and the light chains of rat IgM; cross-adsorbed against mouse immunoglobulins and pooled sera for minimal reactivity
Application Info	Direct and indirect immunofluorescent staining, Enzyme-Linked-Immunosorbent-Assay (ELISA), Fluorescent- Linked-Immunosorbent-Assay (FLISA), Immunoblotting, Immunohistochemistry
Immunogen	Rat IgG
Warning	Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copiousamounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoicacid under acidic conditions. Dilute azide- containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

Image 1       Image 1         Image 1       Image 1	Dilution	Immunofluorescence: FLISA: ELISA: FITC and BIOT conjugates R-PE conjugate TRITC and TXRD conjugates AP conjugate HRP conjugate BGAL conjugate BIOT conjugate
Characterization       To ensure lot- to- lot consistency, each batch of product is tested by ELISA, FLISA, and/or flow cytometry forconformance to characteristics of a standard reference reagent.         Purification       Affinity chromatography on pooled IgG covalently linked to agarose.         Tested Species Reactivity       Rat         Application       FC, IHC-F, IHC-P, ICC, WB         Image 1       Image 1		<= 0.1 ug/10 <sup>6</sup> cells 1:200-1:400 1:2,000-1:4,000 1:4,000-1:8,000 1:500 1:5,000-1:20,000
Characterization       forconformance to characteristics of a standard reference reagent.         Purification       Affinity chromatography on pooled IgG covalently linked to agarose.         Tested Species       Rat         Application       FC, IHC-F, IHC-P, ICC, WB         Image 1       Image 1	Cross Absorption	Pooled mouse sera and pooled mouse plasmacytoma/hybridoma proteins.
Tested Species Reactivity       Rat         Application       FC, IHC-F, IHC-P, ICC, WB         Image 1       Image 1	Characterization	
Reactivity     Rat       Application     FC, IHC-F, IHC-P, ICC, WB       Image 1     Image 1	Purification	Affinity chromatography on pooled IgG covalently linked to agarose.
Image 1		Rat
Image 1	Application	FC, IHC-F, IHC-P, ICC, WB
	Image 1 AVIVA SYSTEMS BIOLO	GY manufactures and sells quality antibody products covering genome wide proteins.
This product is for Research Use Only. Not for diagnostic, human, or veterinary use.		
Optimal conditions of its use should be determined by end users.	Optimal conditions of its use	should be determined by end users.

AVIVA SYSTEMS BIOLOGY 6370 Nancy Ridge Dr., Suite 104, San Diego, CA 92121 USA | Tel: (858)552-6979 | info@avivasysbio.com