



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0


F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

<b>Product Number</b>	OASB02123
<b>Product Page</b>	<a href="http://www.avivasysbio.com/rabbit-anti-goat-igg-h-l-human-sp-ads-antibody-oasb02123.html">www.avivasysbio.com/rabbit-anti-goat-igg-h-l-human-sp-ads-antibody-oasb02123.html</a>
<b>Name</b>	Rabbit Anti-Goat IgG(H+L) Antibody (OASB02123)
<b>Isotype</b>	IgG
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1.0 mg/mL
<b>Gene Full Name</b>	Ig gamma-1 chain C region
<b>Reconstitution and Storage</b>	Store at 2-8C
<b>Storage</b>	<ul style="list-style-type: none"> <li>- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of 100 mM boratebuffered saline, pH 8.2. No preservatives or amine- containing buffer salts added. Store at 2- 8 C</li> <li>- The fluorescein (FITC) conjugate is supplied as 1.0 mg in 1.0 mL PBS/NaN<sub>3</sub>. Store at 2- 8 C</li> <li>- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL of stock solution in 50mM Tris/1mMMgCl<sub>2</sub>/50% Glycerol, pH 8.0, containing 0.1% NaN<sub>3</sub> as preservative. Store at 2- 8 C or long- term at - 20 C</li> <li>- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50%PBS, pH 7.4. No preservative added. Store at 2- 8 C or long- term at - 20 C</li> <li>- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL PBS/NaN<sub>3</sub>. Store at 2- 8 C</li> <li>- Protect fluorochrome- conjugated forms from light. Reagents are stable for the period shown on the label ifstored as directed.</li> </ul>
<b>Other Applications Data</b>	Since applications vary, you should determine the optimum workingdilution of the product that is appropriate for your specific need.
<b>Datasheets/Manuals</b>	Printable datasheet for <a href="#">anti-IGHG1 (OASB02123) antibody</a>
<b>Specificity</b>	IgG(H+L)
<b>Additional Information</b>	<b>Description:</b> Pooled antisera from rabbits hyperimmunized with goat IgG; purified by affinity chromatography on goat IgG covalently linked to agarose; reacts with the heavy and light chains of goat IgG; cross-adsorbed against human serum proteins (SP) for minimal reactivity
<b>Application Info</b>	Immunofluorescent staining, Enzyme-Linked-Immunsorbent-Assay (ELISA), Fluorescent-Linked-Immunsorbent-Assay (FLISA), Western blotting, Dot-and slot-immunoblotting, Immunohistochemistry
<b>Immunogen</b>	Goat IgG
<b>Warning</b>	Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin,eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash withcopious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxichydrazoic acid under acidic conditions. Dilute azide- containing compounds in running water before discardingto avoid accumulation of potentially explosive deposits in lead or copper plumbing.
<b>Dilution</b>	FLISA: ELISA: FITC conjugate AP conjugate HRP conjugate BIOT conjugate 1:200 1:400 1:2,000-1:4,000 1:4,000-1:8,000 1:5,000-1:20,000
<b>Cross Absorption</b>	Pooled human serum proteins.

<b>Characterization</b>	To ensure lot- to- lot consistency, each batch of product is tested by ELISA or FLISA for conformance to characteristics of a standard reference reagent.
<b>Purification</b>	Affinity chromatography on goat IgG covalently coupled to agarose.
<b>Tested Species Reactivity</b>	Goat
<b>Application</b>	FC, IHC-F, IHC-P, ICC, WB
<b>Image 1</b>	 A schematic diagram of a Y-shaped antibody molecule. It consists of two heavy chains (inner lines) and two light chains (outer lines) joined at their C-termini. The two antigen-binding sites are formed by the variable regions of the light and heavy chains, which are shown as the upper arms of the Y.

AVIVA SYSTEMS BIOLOGY manufactures and sells quality antibody products covering genome wide proteins.

This product is for Research Use Only. Not for diagnostic, human, or veterinary use.  
Optimal conditions of its use should be determined by end users.

---

AVIVA SYSTEMS BIOLOGY  
6370 Nancy Ridge Dr., Suite 104, San Diego, CA 92121 USA | Tel: (858)552-6979 | [info@avivasysbio.com](mailto:info@avivasysbio.com)