



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Rat IgG-heavy and light chain Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Catalog No. A110-122A

Lot No. A110-122A-1



|                              |  |
|------------------------------|--|
| <b>APPLICATIONS</b>          | WB, IHC, ICC, ELISA  |
| <b>SPECIES REACTIVITY</b>    | Rat  |
| <b>ISOTYPE</b>               | IgG  |
| <b>AMOUNT</b>                | 1 ml at 1 mg/ml  |
| <b>STORAGE/SHELF LIFE</b>    | 2 - 8° C / 2 years from date of receipt  |
| <b>PHYSICAL STATE</b>        | Liquid   |
| <b>BUFFER</b>                | Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide  |
| <b>ORIGIN</b>                | USA  |
| <b>PRODUCTION PROCEDURES</b> | The antibody was isolated by affinity chromatography using antigen coupled to agarose beads.<br><br>Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.   |
| <b>APPLICATIONS</b>          | By immunoelectrophoresis and ELISA this antibody reacts specifically with rat IgG and with light chains common to other rat immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species.<br><br>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.<br><br>Western Blot                      1:1,000 - 1:30,000<br><br>Immunohistochemistry      1:200 - 1:2,000<br><br>Immunocytochemistry      1:200 - 1:2,000<br><br>ELISA                              1:1,000 - 1:30,000; for coating plates 1:100 - 1:500 |
| <b>APPLICATION NOTES</b>     | Not all listed applications have been specifically tested by our laboratory.   |
| <b>ADDITIONAL INFO</b>       | <a href="http://www.bethyl.com/product/A110-122A">http://www.bethyl.com/product/A110-122A</a><br><br>Use the link above to view SDS, a current list of citations, and other product specific information.  |