

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



#### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# Rat IgG Heavy and Light Chain Cross-Adsorbed Antibody



Donkey Polyclonal Conjugate HRP

Antigen Affinity Purified
Catalog No. A110-337P
Lot No. A110-337P-15

**APPLICATIONS** WB, IHC, ICC, ELISA

SPECIES REACTIVITY Rat. Minimum reactivity to bovine, chicken, goat, human, mouse, rabbit and sheep

AMOUNT 1 ml

CONCENTRATION 0.5 mg/ml

**STORAGE/SHELF LIFE**  $2 - 8^{\circ}\text{C} / 1$  year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.1% Pro-Clean 400

ISOTYPE IgG
ORIGIN USA

**PRODUCTION** Antiserum was cross adsorbed using bovine, human, mouse, rabbit & sheep

PROCEDURES immunosorbents to remove cross reactive antibodies. The antibody to rat IgG was isolated

by affinity chromatography using antigen coupled to agarose beads and conjugated to

horseradish peroxidase (HRP).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of

1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with rat IgG and with light chains common to other rat immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to bovine, human, mouse, rabbit & sheep IgG was detected. This antibody may cross react with IgG from other species.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:5,000 - 1:50,000

Immunohistochemistry 1:200 – 1:5,000

Immunocytochemistry 1:200 - 1:5,000

ELISA 1:10,000 - 1:100,000

**APPLICATION NOTES** Not all listed applications have been specifically tested by our laboratory.

ADDITIONAL INFO https://www.fortislife.com/p/A110-337P

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: December 5, 2022