



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Rabbit IgG-heavy and light chain Antibody

F(ab')<sub>2</sub> Goat Polyclonal                      Conjugate    FITC  
Antigen Affinity Purified  
Catalog No.    A120-114F  
Lot No.            A120-114F-4



---

**APPLICATIONS**                      IHC, ICC, F, IF  
**SPECIES REACTIVITY**              Rabbit  
**ISOTYPE**                              IgG  
**AMOUNT**                              1 ml at 0.5 mg/ml  
**STORAGE/SHELF LIFE**              2 - 8° C / 1 year from date of receipt  
**PHYSICAL STATE**                    Liquid  
**FLUOROPHORE/PROTEIN**          6.1  
**BUFFER**                              Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide  
**ORIGIN**                                USA  
**PRODUCTION PROCEDURES**        The antibody was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')<sub>2</sub> fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to fluorescein isothiocyanate (FITC).

Antibody concentration was determined by extinction coefficient prior to conjugation: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

**APPLICATIONS**                      Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry    1:50 - 1:500  
Immunocytochemistry    1:50 - 1:500  
Flow Cytometry            1:50 - 1:200  
Immunofluorescence    1:50 - 1:500

**APPLICATION NOTES**              Not all listed applications have been specifically tested by our laboratory.

**ADDITIONAL INFO**                    <https://www.bethyl.com/product/A120-114F>  
Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Eric McIntush, PhD | Chief Scientific Officer

Date: December 3, 2018