

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

Rabbit IgG-Fc Fragment Antibody

F(ab')2 Goat Polyclonal Antigen Affinity Purified Catalog No. A120-115A

Lot No. A120–115A–1



APPLICATIONS WB, IHC, ICC, ELISA SPECIES REACTIVITY Rabbit ISOTYPE IgG AMOUNT 1 ml at 0.5 mg/ml STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt PHYSICAL STATE Liquid BUFFER Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide ORIGIN USA PRODUCTION Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab)2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species. APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 - 1:30,000 Immunohistochemistry 1:200 - 1:2,000 Immunohistochemistry 1:200 - 1:30,000; for coating plates 1:100 - 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory. ADDIT		LABORATORIES, IN
ISOTYPE IgG AMOUNT 1 ml at 0.5 mg/ml STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt PHYSICAL STATE Liquid BUFFER Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide ORIGIN USA PRODUCTION Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species. APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 - 1:2,000 Immunohistochemistry 1:200 - 1:2,000 Immunocytochemistry 1:200 - 1:30,000; for coating plates 1:100 - 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.	APPLICATIONS	WB, IHC, ICC, ELISA
AMOUNT I ml at 0.5 mg/ml STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt PHYSICAL STATE Liquid BUFFER Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide ORIGIN USA PROCEDURES Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgC. By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgC. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgC from other species. APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 - 1:30,000 Immunohistochemistry 1:200 - 1:2,000 ELISA 1:1,000 - 1:30,000; for coating plates 1:100 - 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.	SPECIES REACTIVITY	Rabbit
STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt PHYSICAL STATE Liquid BUFFER Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide ORIGIN USA PRODUCTION PROCEDURES Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species. APPLICATIONS Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 – 1:30,000 Immunohistochemistry 1:200 – 1:2,000 Immunocytochemistry 1:200 – 1:2,000 ELISA 1:1,000 – 1:30,000; for coating plates 1:100 – 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.	ISOTYPE	lgG
PHYSICAL STATE Liquid BUFFER Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide ORIGIN USA PRODUCTION PROCEDURES Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species. APPLICATIONS Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 – 1:30,000 Immunocytochemistry 1:200 – 1:2,000 LISA 1:1,000 – 1:30,000; for coating plates 1:100 – 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.	AMOUNT	1 ml at 0.5 mg/ml
BUFFER Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide ORIGIN USA PRODUCTION PROCEDURES Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species. APPLICATIONS Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 – 1:30,000 Immunohistochemistry 1:200 – 1:2,000 ELISA 1:1,000 – 1:30,000; for coating plates 1:100 – 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.	STORAGE/SHELF LIFE	2 - 8° C / 1 year from date of receipt
ORIGIN USA PRODUCTION PROCEDURES Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species. APPLICATIONS Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 – 1:30,000 Immunohistochemistry 1:200 – 1:2,000 ELISA APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.	PHYSICAL STATE	Liquid
PRODUCTION PROCEDURES Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species. APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 – 1:30,000 Immunohistochemistry 1:200 – 1:2,000 ELISA 1:1,000 – 1:30,000; for coating plates 1:100 – 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.	BUFFER	Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide
PROCEDURES isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species. APPLICATIONS Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 – 1:30,000 Immunohistochemistry 1:200 – 1:2,000 ELISA 1:1,000 – 1:30,000; for coating plates 1:100 – 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.	ORIGIN	USA
equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species. APPLICATIONS Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1,000 – 1:30,000 Immunohistochemistry 1:200 – 1:2,000 ELISA 1:1,000 – 1:30,000; for coating plates 1:100 – 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.		isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been
APPLICATIONSCentrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.Western Blot1:1,000 - 1:30,000Immunohistochemistry1:200 - 1:2,000Immunocytochemistry1:200 - 1:30,000; for coating plates 1:100 - 1:500APPLICATION NOTESNot all listed applications have been specifically tested by our laboratory.		
experimentally by the investigator. Prepare working dilution immediately before use.Western Blot1:1,000 – 1:30,000Immunohistochemistry1:200 – 1:2,000Immunocytochemistry1:200 – 1:2,000ELISA1:1,000 – 1:30,000; for coating plates 1:100 – 1:500APPLICATION NOTESNot all listed applications have been specifically tested by our laboratory.ADDITIONAL INFOhttps://www.bethyl.com/product/A120–115A		reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin
Immunohistochemistry 1:200 - 1:2,000 Immunocytochemistry 1:200 - 1:2,000 ELISA 1:1,000 - 1:30,000; for coating plates 1:100 - 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory. ADDITIONAL INFO https://www.bethyl.com/product/A120-115A	APPLICATIONS	
Immunocytochemistry1:200 - 1:2,000ELISA1:1,000 - 1:30,000; for coating plates 1:100 - 1:500APPLICATION NOTESNot all listed applications have been specifically tested by our laboratory.ADDITIONAL INFOhttps://www.bethyl.com/product/A120-115A		Western Blot 1:1,000 – 1:30,000
ELISA 1:1,000 – 1:30,000; for coating plates 1:100 – 1:500 APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory. ADDITIONAL INFO https://www.bethyl.com/product/A120–115A		Immunohistochemistry 1:200 – 1:2,000
APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory. ADDITIONAL INFO https://www.bethyl.com/product/A120-115A		Immunocytochemistry 1:200 – 1:2,000
ADDITIONAL INFO https://www.bethyl.com/product/A120–115A		ELISA 1:1,000 - 1:30,000; for coating plates 1:100 - 1:500
	APPLICATION NOTES	Not all listed applications have been specifically tested by our laboratory.
	ADDITIONAL INFO	

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.Eric McIntush, PhD | Chief Scientific OfficerDate: December 3, 2018

Bethyl Laboratories, Inc. • 25043 West FM 1097 • Montgomery, TX 77356 • 800.338.9579 • 936.597.6111 • 866.597.6105 (FAX) • www.bethyl.com • technical@bethyl.com