

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

## Rabbit IgG-Fc Fragment Antibody

Conjugate Biotin

F(ab')2 Goat Polyclonal Antigen Affinity Purified Catalog No. A120-115B Lot No. A120-115B-1



APPLICATIONS       WB, IHC, ICC, ELISA         SPECIES REACTIVITY       Rabbit         ISOTYPE       IgC         AMOUNT       1 ml at 0.5 mg/ml         STORAGE/SHELF LIFE       2 - 8° C / 1 year from date of receipt         PHSICAL STATE       Liquid         BUFFER       Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide         ORIGIN       USA         PROCEDURES       Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab)2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to biolin.         Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.         By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross react with IgG from other species.         AppLICATIONS       Centrifuge tube to remove product from Id. Optimal working dilution should be determined experimentally by the investigator. Prepare working dilution immediately before use.         Western Blot       1:10.000 – 1:200.000         Immunohistochemistry       1:20 – 1:25,00         Immunohistochemistry       1:10.001 – 1:200.000         RAPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.         ADDITIONAL INFO	LUT NO.		LABORATORIES, INC	
SOTYPEIgGAMOUNT1 ml at 0.5 mg/mlSTORAGE/SHELF LIFE2 - 8° C / 1 year from date of receiptPHYSICAL STATELiquidBUFFERPhosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium AzideORIGINUSAPRODUCTIONAntiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to biotin.Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.AppLICATIONSCentrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution should be determined experimentally by the investigator. Prepare working dilution should be determined experimentally by the investigator. Prepare working dilution immediately before use.Western Blot1:10,000 - 1:200,000Immunocytochemistry1:10,001 - 1:200,000LISA1:10,000 - 1:200,000AppLICATION NOTESNot all listed applications have been specifically tested by our laboratory.	APPLICATIONS	WB, IHC, ICC, ELISA		
AMOUNT       1 ml at 0.5 mg/ml         STORAGE/SHELF LIFE       2 - 8° C / 1 year from date of receipt         PHYSICAL STATE       Liquid         BUFFER       Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide         ORIGIN       USA         PRODUCTION       Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab)2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to biotin.         Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.         By immunoelectrophoresis and ELISA this antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species.         APPLICATIONS       Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.         Western Blot       1:10,000 - 1:200,000         Immunohistochemistry       1:210,010 - 1:500         ELISA       1:10,000 - 1:200,000         APPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.	SPECIES REACTIV	ITY Rabbit		
STORAGE/SHELF LIFE       2 - 8° C / 1 year from date of receipt         PHYSICAL STATE       Liquid         BUFFER       Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide         ORIGIN       USA         PRODUCTION PROCEDURES       Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab)2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to biotin.         Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.         By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species.         APPLICATIONS       Centrifuge tube to remove product from lid. Optimal working dilution immediately before use.         Western Blot       1:10,000 – 1:200,000         Immunohistochemistry       1:250 – 1:2,500         Immunocytochemistry       1:100 – 1:200,000         APPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.         ADDITIONAL INFO       https://www.bethyl.com/product/A120-115B	ISOTYPE	IgG		
PHYSICAL STATE       Liquid         BUFFER       Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide         ORIGIN       USA         PRODUCTION PROCEDURES       Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to biotin.         Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.         By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species.         APPLICATIONS       Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.         Western Blot       1:10,000 – 1:200,000         Immunocytochemistry       1:250 – 1:2,500         Immunocytochemistry       1:10,000 – 1:200,000         APPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.         ADDITIONAL INFO       https://www.bethyl.com/product/A120–115B	AMOUNT	1 ml at 0.5 mg/ml		
BUFFER       Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide         ORIGIN       USA         PRODUCTION PROCEDURES       Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to biotin.         Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.         By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species.         APPLICATIONS       Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.         Western Blot       1:10,000 – 1:200,000         Immunohistochemistry       1:250 – 1:2,500         Immunocytochemistry       1:10,000 – 1:200,000         APPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.         ADDITIONAL INFO       https://www.bethyl.com/product/A120-115B	STORAGE/SHELF	LIFE 2 - 8° C / 1 year from date of receipt		
ORIGIN       USA         PRODUCTION PROCEDURES       Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to biotin.         Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.         By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species.         APPLICATIONS       Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.         Western Blot       1:10,000 – 1:200,000         Immunohistochemistry       1:250 – 1:2,500         Immunocytochemistry       1:10,000 – 1:200,000         MAPPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.	PHYSICAL STATE	Liquid		
PRODUCTION PROCEDURES       Antiserum was solid phase adsorbed to ensure class specificity. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab)2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to biotin.         Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.         By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgC from other species.         APPLICATIONS       Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.         Western Blot       1:10,000 - 1:200,000 Immunohistochemistry         Immunocytochemistry       1:250 - 1:2,500 Immunocytochemistry         ELISA       1:10,000 - 1:200,000         APPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.	BUFFER	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide		
PROCEDURES       isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to biotin.         Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.         By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species.         APPLICATIONS       Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.         Western Blot       1:10,000 - 1:200,000         Immunocytochemistry       1:250 - 1:2,500         Immunocytochemistry       1:10,000 - 1:200,000         APPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.	ORIGIN	USA		
equals 1.0 mg of IgG.         By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG. Cross reactivity with IgA and IgM is negligible. No antibody was detected against non-immunoglobulin serum proteins. This antibody may cross react with IgG from other species.         APPLICATIONS       Centrifuge tube to remove product from Iid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.         Western Blot       1:10,000 - 1:200,000         Immunohistochemistry       1:250 - 1:2,500         Immunocytochemistry       1:10,000 - 1:200,000         APPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.         ADDITIONAL INFO       https://www.bethyl.com/product/A120-115B		isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been		
APPLICATIONSCentrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.Western Blot1:10,000 - 1:200,000Immunohistochemistry1:250 - 1:2,500Immunocytochemistry1:100 - 1:200,000ELISA1:10,000 - 1:200,000Not all listed applications have been specifically tested by our laboratory.ADDITIONAL INFOhttps://www.bethyl.com/product/A120-115B			coefficient: absorbance at 280 nm of 1.4	
experimentally by the investigator. Prepare working dilution immediately before use.Western Blot1:10,000 - 1:200,000Immunohistochemistry1:250 - 1:2,500Immunocytochemistry1:100 - 1:500ELISA1:10,000 - 1:200,000APPLICATION NOTESNot all listed applications have been specifically tested by our laboratory.ADDITIONAL INFOhttps://www.bethyl.com/product/A120-115B	reactivity with IgA and IgM is negligible. No antibody was detected		vas detected against non-immunoglobulin	
Immunohistochemistry       1:250 - 1:2,500         Immunocytochemistry       1:100 - 1:500         ELISA       1:10,000 - 1:200,000         APPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.         ADDITIONAL INFO       https://www.bethyl.com/product/A120-115B	APPLICATIONS			
Immunocytochemistry       1:100 - 1:500         ELISA       1:10,000 - 1:200,000         APPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.         ADDITIONAL INFO       https://www.bethyl.com/product/A120-115B		Western Blot 1:10,000 - 1:200,000		
APPLICATION NOTES       ELISA       1:10,000 - 1:200,000         Not all listed applications have been specifically tested by our laboratory.         ADDITIONAL INFO       https://www.bethyl.com/product/A120-115B		Immunohistochemistry 1:250 – 1:2,500		
APPLICATION NOTES       Not all listed applications have been specifically tested by our laboratory.         ADDITIONAL INFO       https://www.bethyl.com/product/A120-115B		Immunocytochemistry 1:100 – 1:500		
ADDITIONAL INFO https://www.bethyl.com/product/A120–115B		ELISA 1:10,000 - 1:200,000		
	APPLICATION NO	<b>TES</b> Not all listed applications have been specifically tested	by our laboratory.	
	ADDITIONAL INFO			

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.Eric McIntush, PhD | Chief Scientific OfficerDate: December 3, 2018

Bethyl Laboratories, Inc. • 25043 West FM 1097 • Montgomery, TX 77356 • 800.338.9579 • 936.597.6111 • 866.597.6105 (FAX) • www.bethyl.com • technical@bethyl.com