

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



## Rabbit IgG-heavy and light chain cross-adsorbed Antibody

F(ab')2 Goat Polyclonal Conjugate HRP

Antigen Affinity Purified

Catalog No. A120–314P

Lot No. A120–314P–5



**APPLICATIONS** WB, IHC, ICC, ELISA

SPECIES REACTIVITY Rabbit. Minimum reactivity to bovine, chicken, horse, human, mouse, rat and sheep

AMOUNT 1 ml

CONCENTRATION 0.5 mg/ml

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.05% Pro-Clean 400

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antiserum was cross adsorbed using bovine, chicken, horse, human, mouse, rat and sheep immunosorbents to remove cross reactive antibodies. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to horseradish peroxidase (HRP).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG and with light chains common to other rabbit immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to bovine, chicken, horse, human, mouse, rat and sheep IgG was detected. This antibody may cross react with IgG from other

species.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:5,000 – 1:50,000

Immunohistochemistry 1:200 - 1:5.000

Immunocytochemistry 1:200 - 1:5,000

ELISA 1:10,000 - 1:100,000

**APPLICATION NOTES** Not all listed applications have been specifically tested by our laboratory.

ADDITIONAL INFO https://www.bethyl.com/product/A120-314P

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Eric McIntush, PhD | Chief Scientific Officer

Date: June 18, 2019

314P