



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Rabbit IgG-heavy and light chain human-adsorbed Antibody

Goat Polyclonal

Antigen Affinity Purified

Catalog No. A120-401A

Lot No. A120-401A-1



<b>APPLICATIONS</b>	WB, IHC, ICC, ELISA								
<b>SPECIES REACTIVITY</b>	Rabbit. Minimum reactivity to human								
<b>ISOTYPE</b>	IgG								
<b>AMOUNT</b>	1 ml at 0.5 mg/ml								
<b>STORAGE/SHELF LIFE</b>	2 - 8° C / 1 year from date of receipt								
<b>PHYSICAL STATE</b>	Liquid								
<b>BUFFER</b>	Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide								
<b>ORIGIN</b>	USA								
<b>PRODUCTION PROCEDURES</b>	<p>Antiserum was cross adsorbed using a human immunosorbent to remove cross reactive antibodies. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads.</p> <p>Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p> <p>By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG and with light chains common to other rabbit immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 0.1% cross reactivity to human IgG was detected. This antibody may cross react with IgG from other species.</p>								
<b>APPLICATIONS</b>	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Western Blot</td><td>1:1,000 - 1:20,000</td></tr><tr><td>Immunohistochemistry</td><td>1:100 - 1:1,000</td></tr><tr><td>Immunocytochemistry</td><td>1:100 - 1:1,000</td></tr><tr><td>ELISA</td><td>1:1,000 - 1:20,000; for coating plates 1:50 - 1:250</td></tr></table>	Western Blot	1:1,000 - 1:20,000	Immunohistochemistry	1:100 - 1:1,000	Immunocytochemistry	1:100 - 1:1,000	ELISA	1:1,000 - 1:20,000; for coating plates 1:50 - 1:250
Western Blot	1:1,000 - 1:20,000								
Immunohistochemistry	1:100 - 1:1,000								
Immunocytochemistry	1:100 - 1:1,000								
ELISA	1:1,000 - 1:20,000; for coating plates 1:50 - 1:250								
<b>APPLICATION NOTES</b>	Not all listed applications have been specifically tested by our laboratory.								
<b>ADDITIONAL INFO</b>	<p><a href="https://www.bethyl.com/product/A120-401A">https://www.bethyl.com/product/A120-401A</a> Use the link above to view SDS, a current list of citations, and other product specific information.</p>								

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Eric McIntush, PhD | Chief Scientific Officer

Date: December 3, 2018