



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Monkey IgG Heavy and Light Chain Antibody Cross-Adsorbed

Goat Polyclonal Conjugate HRP

Antigen Affinity Purified

Catalog No. A140-202P

Lot No. A140-202P-19

**APPLICATIONS** WB, IHC, ICC, ELISA

**SPECIES REACTIVITY** Monkey. Minimum reactivity to human

**AMOUNT** 1 ml

**CONCENTRATION** 0.5 mg/ml

**STORAGE/SHELF LIFE** 2 – 8°C / 1 year from date of receipt

**PHYSICAL STATE** Liquid

**BUFFER** Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.05% Pro-Clean 400

**ISOTYPE** IgG

**ORIGIN** USA

**PRODUCTION PROCEDURES** The antibody was adsorbed against human serum, isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to horseradish peroxidase (HRP).

Antibody concentration was determined by extinction coefficient prior to conjugation: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. Molar enzyme/antibody protein ratio is 4:1.

By immunoelectrophoresis and ELISA this antibody reacts specifically with monkey IgG and with light chains common to other monkey immunoglobulins. Species detected are Rhesus, Baboon, Carb Green, Cyno, Chimp and African Green. Reaction to human IgG was less than 1%. No antibody was detected against non-immunoglobulin serum proteins.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000 – 1:30,000

Immunohistochemistry 1:200 – 1:500

Immunocytochemistry 1:200 – 1:500

ELISA 1:10,000 – 1:100,000

**APPLICATION NOTES** Not all listed applications have been specifically tested by our laboratory.

**ADDITIONAL INFO** <https://www.fortislife.com/p/A140-202P>

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Michael Spencer, PhD Date: October 31, 2022