

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



## **EGFR Antibody**

Rabbit Polyclonal

Antigen Affinity Purified Protein ID P00533

Catalog No. A300–387A–T GeneID 1956

Lot No. A300-387A-T-1

APPLICATIONS WB, IP, IHC
SPECIES REACTIVITY Human

**PRESUMED REACTIVITY** Based on 100% sequence identity, this antibody is predicted to react with Monkey

AMOUNT 20 μl (2 blots)

CONCENTRATION 20 μg/ml

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to EGFR immobilized on solid support.

The epitope recognized by A300-387A-T maps to a region between residues 1100 and 1150 of human Epidermal Growth Factor Receptor using the numbering given in Swiss-Prot entry P00533

(GeneID 1956).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1000

Immunoprecipitation The antibody contained within A300–387A–T has been qualified for

use in immunoprecipitation; however, we recommend using the alternative formulation of this antibody found as product A300–387A.

Immunohistochemistry 1:10 - 1:40. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE tissue sections.

**APPLICATION NOTES** Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.


ADDITIONAL INFO https://www.bethyl.com/product/A300-387A-T

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019

EGFR Antibody A300-387A-T



Detection of human EGFR by western blot. Samples: Whole cell lysate (50  $\mu$ g) from HeLa and MCF-7 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-EGFR antibody A300-387A-T (lot A300-387A-T-1) used at 1:1000. Detection: Chemiluminescence with an exposure time of 30 seconds.