



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien



T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Detection of human and mouse SERCA2 by western blot.
Samples: Whole cell lysate (50 µg) from HeLa, HEK293T, and mouse NIH 3T3 cells prepared using NETN lysis buffer.
Antibody: Affinity purified rabbit anti-SERCA2 antibody A300-406A (lot A300-406A-4) used for WB at 0.1 µg/ml.
Detection: Chemiluminescence with an exposure time of 30 seconds.

Detection of human SERCA2 by western blot of immunoprecipitates.
Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HEK293T cells prepared using NETN lysis buffer.
Antibodies: Affinity purified rabbit anti-SERCA2 antibody A300-406A (lot A300-406A-4) used for IP at 6 µg per reaction. SERCA2 was also immunoprecipitated by a previous lot of this antibody A300-406A-3. For blotting immunoprecipitated SERCA2, A300-406A was used at 1 µg/ml.
Detection: Chemiluminescence with an exposure time of 3 minutes.