

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com


CTCF Antibody

| Rabbit Polyclonal Protein ID NP_006556.1 Catalog No. A300-542A-T GeneID 10664 Lot No. A300-542A-T-1 Image: Catalog No. A300-542A-T-1 APPLICATIONS IP Image: Catalog No. A300-542A-T-1 APPLICATIONS IP Image: Catalog No. Second No. SPECIES REACTIVITY Human Image: Catalog No. MOUNT Image: Catalog No. PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Rat AMOUNT Image: Catalog No. CONCENTRATION 10 µl Image: Catalog No. Second No. Second No. STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt PHYSICAL STATE Liquid BUFFER IgG Image: Catalog No. Sold No. ORIGIN USA Image: Catalog No. Sold No. PRODUCTION Matibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664). Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. | | ay | | | | |
|--|--------------------|-------------------|--|---------------------------|---|--|
| Lot No. A300-542A-T-1 APPLICATIONS IP SPECIES REACTIVITY Human PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Rat AMOUNT 10 μl CONCENTRATION 1000 μg/ml STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt PHYSICAL STATE Liquid BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide ISOTYPE IgG ORIGIN USA PRODUCTION Antibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664). Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot Not recommended. Use rabbit anti-CTCF antibody A300-543A-T. Immunoprecipitation 2 - 5 µg/mg lysate APPLICATION NOTES Western Blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Coat | • | | Protein ID | NP_006556.1 | | |
| APPLICATIONS IP SPECIES REACTIVITY Human PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Rat AMOUNT 10 µl CONCENTRATION 1000 µg/ml STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt PHYSICAL STATE Liquid BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide ISOTYPE IgG ORIGIN USA PRODUCTION PROCEDURES Antibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664). Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgC. APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot Not recommended. Use rabbit anti-CTCF antibody A300-543A-T. Immunoprecipitation 2 - 5 µg/mg lysate APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Coat anti-R | Catalog No. A3 | 00-542A-T | GenelD | 10664 | | |
| SPECIES REACTIVITY Human RESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Rat AMOUNT 10 µl CONCENTRATION 100 µg/ml STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt PHYSICAL STATE Liquid BUFFER 10 G ORIGIN USA PRODUCTION PROCEDURES Antibody was affinity purified using an epitope specific to CTCF immobilized on solid support. Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 m of lgC. APPLICATIONS Centrifuge tube to remover product from lid. Optimal working dilution should be determined experimentally by the investigator. Prepare working dilution immediately before use. APPLICATION NOTES Vestern Blot Not recommended. Use rabbit anti-CTCF antibody A300-543A-T. APPLICATION NOTES Centrifuge tube to remover product from lid. Optimal working dilution should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot Not recommended. Use rabbit anti-CTCF antibody A300-543A-T. Immunoprecipitation 2 - 5 µg/mg lysate APPLICATION NOTES Western Blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Cat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (Link to IP-western Blot views USA, a current list of citations, and other product specific information. | Lot No. A3 | 00-542A-T-1 | | | | |
| PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Rat AMOUNT 10 μl CONCENTRATION 1000 µg/ml STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt PHYSICAL STATE Liquid BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide ISOTYPE IgG ORICIN USA PRODUCTION Antibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664). Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot Not recommended. Use rabbit anti-CTCF antibody A300-543A-T. Immunoprecipitation 2 - 5 µg/mg lysate APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below). ADDITIONAL INFO <t< th=""><th>APPLICATIONS</th><th>IP</th><th></th><th></th><th></th></t<> | APPLICATIONS | IP | | | | |
| AMOUNT10 μlCONCENTRATION1000 μg/mlSTORAGE/SHELF LIFE2 - 8° C / 1 year from date of receiptPHYSICAL STATELiquidBUFFERTris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium AzideISOTYPEIgGORIGINUSAPRODUCTION PROCEDURESAntibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664).APPLICATIONSCentrifuge tube to remove product from lid. Optimal working dilutions should be determined | SPECIES REACTIVITY | Human | Human | | | |
| AMOUNT10 μlCONCENTRATION1000 μg/mlSTORAGE/SHELF LIFE2 - 8° C / 1 year from date of receiptPHYSICAL STATELiquidBUFFERTris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium AzideISOTYPEIgGORIGINUSAPRODUCTION PROCEDURESAntibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664).APPLICATIONSCentrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution should A300-543A-T. Immunoprecipitation 2 - 5 μg/mg lysateAPPLICATION NOTESWestern Blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).ADDITIONAL INFOhttps://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | PRESUMED REACTIV | ITY Based on 1009 | used on 100% sequence identity, this antibody is predicted to react with Rat | | | |
| CONCENTRATION1000 μg/mlSTORAGE/SHELF LIFE2 - 8° C / 1 γear from date of receiptPHYSICAL STATELiquidBUFFERTris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium AzideISOTYPEIgGORIGINUSAPRODUCTION PROCEDURESAntibody was affinity purified using an epitope specific to CTCF immobilized on solid support.The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664).APPLICATIONSCentrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution should be determined experimentally by the investigator. Prepare working dilution should he determined experimentally by the investigator. Prepare working dilution immediately before use.APPLICATION NOTESWestern Blot Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-wester blot product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | | | | | | |
| STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt PHYSICAL STATE Liquid BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide ISOTYPE IgG ORIGIN USA PRODUCTION PROCEDURES Antibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664). Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot Not recommended. Use rabbit anti-CTCF antibody A300-543A-T. Immunoprecipitation 2 - 5 µg/mg Iysate APPLICATION NOTES Western Blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional linfo section below). ADDITIONAL INFO https://www.bethyl.com/product/A300-542A-T | | | μg/ml | | | |
| PHYSICAL STATE Liquid BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide ISOTYPE IgG ORIGIN USA PRODUCTION PROCEDURES Antibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664). APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. APPLICATION NOTES Western Blot Not recommended. Use rabbit anti-CTCF antibody A300-543A-T. Immunoprecipitation APPLICATION NOTES Western Blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below). ADDITIONAL INFO https://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | STORAGE/SHELF LIF | | | | | |
| BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide ISOTYPE IgG ORIGIN USA PRODUCTION PROCEDURES Antibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664). APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Vestern Blot Not recommended. Use rabbit anti-CTCF antibody A300-543A-T. Immunoprecipitation 2 - 5 µg/mg lysate APPLICATION NOTES Western Blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below). ADDITIONAL INFO https://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | | - | | | | |
| ORIGINUSAPRODUCTION PROCEDURESAntibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664).APPLICATIONSCentrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.APPLICATION NOTESWestern BlotNot recommended. Use rabbit anti-CTCF antibody A300-543A-T. Immunoprecipitation 2 - 5 μg/mg lysateAPPLICATION NOTESWestern blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).ADDITIONAL INFOhttps://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | BUFFER | - | /phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide | | | |
| PRODUCTION PROCEDURESAntibody was affinity purified using an epitope specific to CTCF immobilized on solid support. The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664). Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.APPLICATIONSCentrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot Immunoprecipitation 2 - 5 μg/mg lysateAPPLICATION NOTESWestern blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).ADDITIONAL INFOhttps://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | ISOTYPE | IgG | | | | |
| PROCEDURES The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664). Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot Not recommended. Use rabbit anti-CTCF antibody A300-543A-T. Immunoprecipitation 2 - 5 μg/mg lysate APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below). ADDITIONAL INFO https://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | ORIGIN | USA | USA | | | |
| The epitope recognized by A300-542A-T maps to a region between residues 150 and 200 of human CCCTC-binding factor using the numbering given in entry NP_006556.1 (GeneID 10664).Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.APPLICATIONSCentrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.Western BlotNot recommended. Use rabbit anti-CTCF antibody A300-543A-T. Immunoprecipitation 2 - 5 μg/mg lysateAPPLICATION NOTESWestern blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).ADDITIONAL INFOhttps://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | PRODUCTION | Antibody was a | Antibody was affinity purified using an epitope specific to CTCF immobilized on solid support. | | | |
| APPLICATIONSCentrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.Western BlotNot recommended. Use rabbit anti-CTCF antibody A300-543A-T.Immunoprecipitation2 - 5 μg/mg lysateAPPLICATION NOTESWestern blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).ADDITIONAL INFOhttps://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | PROCEDURES | | | | | |
| experimentally by the investigator. Prepare working dilution immediately before use.Western BlotNot recommended. Use rabbit anti-CTCF antibody A300-543A-T.Immunoprecipitation2 - 5 µg/mg lysateAPPLICATION NOTESWestern blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).ADDITIONAL INFOhttps://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | | | | | ction coefficient: absorbance at 280 nm | |
| Immunoprecipitation2 - 5 µg/mg lysateAPPLICATION NOTESWestern blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).ADDITIONAL INFOhttps://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | | | | | | |
| APPLICATION NOTESWestern blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).ADDITIONAL INFOhttps://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | | Western Blot | No | t recommended. Use rabbit | anti-CTCF antibody A300-543A-T. | |
| Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).ADDITIONAL INFOhttps://www.bethyl.com/product/A300-542A-T Use the link above to view SDS, a current list of citations, and other product specific information. | | Immunoprecip | pitation 2 - | - 5 µg/mg lysate | | |
| Use the link above to view SDS, a current list of citations, and other product specific information. | APPLICATION NOTES | Goat anti-Rabb | Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE | | | |
| | | | | | | |
| | | | | | | |

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.

2



Detection of human CTCF by immunoprecipitation. *Samples:* Whole cell lysate (1 mg for IP; 1/4 of IP loaded/lane) from MCF-7 cells. *Antibody:* Affinity purified rabbit anti-CTCF antibody BL2545 (Cat. No. A300-542A-T) used at 3 μ g/mg lysate for IP. Human CTCF was also immunoprecipitated using affinity purified rabbit anti-CTCF antibodies BL2546 (Cat. No. A300-543A) and BL2547 (Cat. No. A300-544A). CTCF was blotted using BL2546 at 0.03 μ g/ml. *Detection:* Chemiluminescence with exposure times of 10 seconds.

Bethyl Laboratories, Inc. • 25043 West FM 1097 • Montgomery, TX 77356 • 800.338.9579 • 936.597.6111 • 866.597.6105 (FAX) • www.bethyl.com • technical@bethyl.com