

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH


Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

SFRS8/SWAP Antibody


Rabbit Polyclonal Antigen Affinity Purified			Protein ID	NP_004583.2	
Catalog No. A300–985A			GenelD	6433	
Lot No.		985A-1	Jeneid	0.135	LABORATORIES, INC
APPLICATIONS		WB, IP, IHC			
SPECIES REACTIVITY		Human			
AMOUNT		100 µl			
CONCENTRATION		200 µg/ml			
STORAGE/SHELF LIFE		2 – 8° C / 1 year from date of receipt			
PHYSICAL STATE		Liquid			
BUFFER		Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide			
ISOTYPE		IgG			
ORIGIN		USA			
PRODUCTION PROCEDURES		Antibody was affinity purified using an epitope specific to SFRS8/SWAP immobilized on solid support.			
		The epitope recognized by A300–985A maps to a region between residue 500 and 550 of human splicing factor, arginine/serine-rich 8 using the numbering given in entry NP_004583.2 (GeneID 6433).			
		Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.			
APPLICATIONS		Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.			
		Western Blot	1:2	,000 - 1:10,000	
		Immunoprecipita	tion 2 -	5 µg/mg lysate	
		Immunohistoche		00 – 1:1,000. Epitope retriev ommended for FFPE tissue se	
APPLICATION NOTES		Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100–020), Goat anti–Rabbit Light Chain HRP Conjugate (Cat. No. A120–113P) and 4–8% SDS–PAGE (link to IP–western blot protocol in Additional Info section below).			
IHC HUMAN CONTROLS		Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE. Breast Carcinoma, Laryngeal Squamous Cell Carcinoma, Ovarian Carcinoma, Prostate Carcinoma			
ADDITIONAL INFO		https://www.bethyl.com/product/A300-985A Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB			

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.



Detection of human SFRS8/SWAP by western blot and immunoprecipitation. *Samples:* Whole cell lysate (5, 15 and 50 µg for WB; 1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti–SFRS8/SWAP antibody A300–985A used for WB at 0.04 µg/ml (A) and 1 µg/ml (B) and used for IP at 3 µg/mg lysate (B). SFRS8/SWAP was also immunoprecipitated by rabbit anti– SFRS8/SWAP antibody BL4340, which recognizes an upstream epitope. *Detection:* Chemiluminescence with exposure times of 3 minutes (A) and 30 seconds (B).



Detection of human SFRS8/SWAP by

immunohistochemistry. *Sample:* FFPE section of human breast carcinoma. *Antibody:* Affinity purified rabbit anti-SFRS8/SWAP (Cat. No. A300-985A Lot1) used at a dilution of 1:200 (1µg/ml). *Detection:* DAB