



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# ATRX Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP\_000480.2

Catalog No. A301-045A


GeneID 546

Lot No. A301-045A-4




|                              |   |              |                    |                     |                     |                      |  |
|------------------------------|---|--------------|--------------------|---------------------|---------------------|----------------------|--|
| <b>APPLICATIONS</b>          | WB, IP, IHC   |              |                    |                     |                     |                      |  |
| <b>SPECIES REACTIVITY</b>    | Human   |              |                    |                     |                     |                      |  |
| <b>PRESUMED REACTIVITY</b>   | Based on 100% sequence identity, this antibody is predicted to react with Chimpanzee  |              |                    |                     |                     |                      |  |
| <b>AMOUNT</b>                | 100 µl  |              |                    |                     |                     |                      |  |
| <b>CONCENTRATION</b>         | 1000 µg/ml  |              |                    |                     |                     |                      |  |
| <b>STORAGE/SHELF LIFE</b>    | 2 - 8° C / 1 year from date of receipt  |              |                    |                     |                     |                      |  |
| <b>PHYSICAL STATE</b>        | Liquid  |              |                    |                     |                     |                      |  |
| <b>BUFFER</b>                | Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide  |              |                    |                     |                     |                      |  |
| <b>ISOTYPE</b>               | IgG   |              |                    |                     |                     |                      |  |
| <b>ORIGIN</b>                | USA   |              |                    |                     |                     |                      |  |
| <b>PRODUCTION PROCEDURES</b> | <p>Antibody was affinity purified using an epitope specific to ATRX immobilized on solid support. The epitope recognized by A301-045A maps to a region between residue 2442 and the C-terminus (residue 2492) of human alpha thalassemia/mental retardation syndrome X-linked using the numbering given in entry NP_000480.2 (GeneID 546).</p> <p>Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p>        |              |                    |                     |                     |                      |  |
| <b>APPLICATIONS</b>          | <p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Western Blot</td><td>1:2,000 – 1:10,000</td></tr><tr><td>Immunoprecipitation</td><td>2 – 10 µg/mg lysate</td></tr><tr><td>Immunohistochemistry</td><td>1:500 to 1:2,000. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.</td></tr></table> | Western Blot | 1:2,000 – 1:10,000 | Immunoprecipitation | 2 – 10 µg/mg lysate | Immunohistochemistry | 1:500 to 1:2,000. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections. |
| Western Blot                 | 1:2,000 – 1:10,000  |              |                    |                     |                     |                      |  |
| Immunoprecipitation          | 2 – 10 µg/mg lysate   |              |                    |                     |                     |                      |  |
| Immunohistochemistry         | 1:500 to 1:2,000. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.  |              |                    |                     |                     |                      |  |
| <b>APPLICATION NOTES</b>     | <p>Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).</p> <p>Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.</p>   |              |                    |                     |                     |                      |  |
| <b>IHC HUMAN CONTROLS</b>    | Breast Carcinoma, Prostate Carcinoma, Skin Basal Cell Carcinoma   |              |                    |                     |                     |                      |  |
| <b>ADDITIONAL INFO</b>       | <p><a href="https://www.bethyl.com/product/A301-045A">https://www.bethyl.com/product/A301-045A</a></p> <p>Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol <a href="https://www.bethyl.com/content/protocol_IP_WB">https://www.bethyl.com/content/protocol_IP_WB</a></p>   |              |                    |                     |                     |                      |  |


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Eric McIntush, PhD | Chief Scientific Officer Date: October 30, 2019



**Detection of human ATRX by western blot.** *Samples:* Whole cell lysate (50 µg) from HeLa, A-549, HEK293T, and MCF-7 cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-ATRX antibody A301-045A (lot A301-045A-4) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.



**Detection of human ATRX by western blot of immunoprecipitates.** *Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-ATRX antibody A301-045A (lot A301-045A-4) used for IP at 6 µg per reaction. ATRX was also immunoprecipitated by a previous lot of this antibody (lot A301-045A-3) and rabbit anti-ATRX antibody BL4515. For blotting immunoprecipitated ATRX, A301-045A was used at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.



**Detection of human ATRX by immunohistochemistry.** *Sample:* FFPE section of human lung cancer. *Antibody:* Affinity purified rabbit anti-ATRX (Cat. No. A301-045A-4) used at a dilution of 1:1,000 (1 µg/ml). *Detection:* DAB