

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

| KIF18A Antibody | | | | | | | | | |
|---|--------------------|---|-------------|--|---|---------|--|--|--|
| Rabbit Polyclonal | | | | | | | | | |
| Antigen Affinity Purified | | | Protein ID | NP_112494.2 | | í • | | | |
| Catalog No. | alog No. A301-079A | | GenelD | 81930 | DETU | v. | | | |
| Lot No. | A301- | 079A-1 | | | | | | | |
| APPLICATIONS | | IP | | | | | | | |
| SPECIES REACTIVITY | | Human | | | | | | | |
| AMOUNT | | 100 µl | | | | | | | |
| CONCENTRATION | | 1000 μg/ml | | | | | | | |
| STORAGE/SHELF LIFE | | 2 - 8° C / 1 year from date of receipt | | | | | | | |
| PHYSICAL STATE | | Liquid | | | | | | | |
| BUFFER | | Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide | | | | | | | |
| ISOTYPE | | IgG | | | | | | | |
| ORIGIN | | USA | | | | | | | |
| PRODUCTION PROCEDURES | | Antibody was affinity purified using an epitope specific to KIF18A immobilized on solid support. | | | | | | | |
| PROCEDURES | | | | | on between residue 650 and 700 of 1 in entry NP_112494.2 (GeneID 819 | | | | |
| | | Immunoglobuli of 1.4 equals 1 | | on was determined by ext | nction coefficient: absorbance at 2 | 80 nm | | | |
| APPLICATIONS | | Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. | | | | | | | |
| | | Western Blot | No | t recommended. Use rabl | it anti-KIF18A antibody A301-080A | ۹. | | | |
| | | Immunoprecip | itation 2 - | - 5 µg/mg lysate | | | | | |
| APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S10 Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAG (link to IP-western blot protocol in Additional Info section below). | | | | A120-113P) and 4-8% SDS-PAGE | 20), | | | | |
| ADDITIONAL INFO | | https://www.bethyl.com/product/A301–079A | | | | | | | |
| | | | | DS, a current list of citatio ps://www.bethyl.com/cor | ns, and other product specific inform tent/protocol_IP_WB | nation. | | | |

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.

| | IF | P/WB | ; | |
|--------------|----|------|---|--------------|
| kDa 460- | | | | |
| 268_ 238- | | | | |
| 171- | | | | |
| 117- | - | - | | 🗲 KIF18A |
| 71- | | 10 | | |
| 55- | | | | |
| 41- | | | | |
| | + | - | - | A301-079A |
| | - | + | - | A301-080A IP |
| | - | - | + | Ctrl IgG |

Detection of human KIF18A by western blot of

immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-KIF18A antibody A301-079A used for IP at 3 μ g/mg lysate. KIF18A was also immunoprecipitated by rabbit anti-KIF18A antibody A301-080A, which recognizes a downstream epitope. For blotting immunoprecipitated KIF18A, A301-080A was used at 1 μ g/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.