



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0


F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 






#### Detection of human Pericentrin-Kendrin by western blot.

*Samples:* Whole cell lysate (10 µg) from HEK293T, HeLa, Hep-G2, and K-562 cells prepared using NETN lysis buffer.

*Antibody:* Affinity purified rabbit anti-Pericentrin-Kendrin antibody (A301-348A lot 4) used for WB at 0.04 µg/ml.


*Detection:* Chemiluminescence with an exposure time of 30 second.



#### Detection of human Pericentrin-Kendrin by western blot of immunoprecipitates.

*Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HEK293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-Pericentrin-Kendrin antibody (A301-348A lot 4) used for IP at 6 µg per reaction.

Pericentrin-Kendrin was also immunoprecipitated by a previous lot of this antibody (A301-348A lot 3) and another rabbit anti-Pericentrin-Kendrin antibody (BL5517). For blotting immunoprecipitated Pericentrin-Kendrin, A301-348A was used at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 1 second.



**Detection of human Pericentrin/Kendrin by immunocytochemistry.** *Sample:* Formaldehyde-fixed asynchronous HeLa cells. *Antibody:* Affinity purified rabbit anti-Pericentrin/Kendrin (Cat. No.A301-348A Lot2) used at a dilution of 1:1,000 (1 µg/ml). *Detection:* Red-fluorescent goat anti-rabbit IgG-heavy and light chain cross-adsorbed Antibody DyLight® 594 Conjugated used at a dilution of 1:100. Counterstain. DAPI (blue) and Phalloidin (green)