

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH


Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

## ATXN2L Antibody

Rabbit Polyclonal	-			and the second second
Antigen Affinity Purified		Protein ID	NP_009176.2	
Catalog No. A301-	369A	GenelD	11273	
Lot No. A301-	369A-1 BEIRYL			
APPLICATIONS	WB, IP			
SPECIES REACTIVITY	Human			
PRESUMED REACTIVITY	Based on 100% sequence identity, this antibody is predicted to react with Mouse			
AMOUNT	100 µl			
CONCENTRATION	200 µg/ml			
STORAGE/SHELF LIFE	2 - 8° C / 1 year from date of receipt			
PHYSICAL STATE	Liquid			
BUFFER	Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide			
ISOTYPE	IgG			
ORIGIN	USA			
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to ATXN2L immobilized o			to ATXN2L immobilized on solid support.
	The epitope recognized by A301-369A maps to a region between residue 150 and 200 of human ataxin 2-like using the numbering given in entry NP_009176.2 (GeneID 11273).			
	Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.			
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.			
	Western Blot	1:2	2,000 - 1:10,000	
	Immunoprecip	itation 2 -	- 5 μg/mg lysate	
APPLICATION NOTES	Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100–020), Goat anti–Rabbit Light Chain HRP Conjugate (Cat. No. A120–113P) and 4–8% SDS–PAGE (link to IP–western blot protocol in Additional Info section below).			
	Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.			
ADDITIONAL INFO	https://www.bethyl.com/product/A301-369A			
	Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB			

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.



Detection of human ATXN2L by western blot and immunoprecipitation. *Samples:* Whole cell lysate from HeLa (5, 15 and 50 µg for WB; 1 mg for IP, 20% of IP loaded) and HEK293T (T; 50 µg) cells. *Antibodies:* Affinity purified rabbit anti-ATXN2L antibody A301-369A used for WB at 0.04 µg/ml (A) and 0.1 µg/ml (B) and used for IP at 3 µg/mg lysate. ATXN2L was also immunoprecipitated by rabbit anti-ATXN2L antibodies A301-370A and A301-371A, which recognize downstream epitopes. *Detection:* Chemiluminescence with exposure times of 30 seconds (A and B).