



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Uroplakin III Monoclonal Antibody [BC17]

Mouse Monoclonal

Purified		RefSeq ID	NP_008884.1
Catalog No.	A500-042A-T	Uniprot ID	O75631
Lot No.	1	GenID	7380

APPLICATIONS	IHC, ICC
SPECIES REACTIVITY	Human
AMOUNT	10 µl (5+ tests)
CONCENTRATION	500 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Phosphate Buffered Saline (PBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide
ISOTYPE	IgG1
CLONE #	BC17
ORIGIN	USA
PRODUCTION PROCEDURES	Monoclonal antibody was purified from cell culture supernatant.

Uroplakin III

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.


Immunocytochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE cell sections.

IHC HUMAN CONTROLS Bladder Carcinoma, RT-4 Cells


ADDITIONAL INFO <https://www.fortislife.com/p/A500-042A-T>

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: April 28, 2023



Detection of human Uroplakin III by immunohistochemistry. *Sample:* FFPE section of bladder carcinoma. *Antibody:* Mouse anti-Uroplakin III monoclonal antibody [BC12] (A500-042A-1). *Secondary:* HRP-conjugated goat anti-mouse IgG (A90-116P).



Detection of human Uroplakin III by immunocytochemistry. *Sample:* FFPE section of RT-4 cells. *Antibody:* Mouse anti-Uroplakin III monoclonal antibody [BC12] (A500-042A-1). *Secondary:* HRP-conjugated goat anti-mouse IgG (A90-116P).