



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


SARS-CoV S-RBD Recombinant Monoclonal Antibody [CR3022-IgG1]

Human Recombinant Monoclonal

Purified		Protein ID	YP_009724390.1
Catalog No.	A700-201	GeneID	43740568
Lot No.	A700-201-1		

APPLICATIONS	ELISA
AMOUNT	100 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 - 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide
ISOTYPE	IgG1
CLONE #	CR3022-IgG1
ORIGIN	USA
PRODUCTION PROCEDURES	Recombinant antibody was purified from cell culture supernatant. The original monoclonal antibody was generated by sequencing peripheral blood lymphocytes of a patient exposed to SARS-CoV.
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. ELISA
ADDITIONAL INFO	https://www.bethyl.com/product/A700-201 Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: May 3, 2022



Analysis of Recombinant SARS-CoV Spike RBD [CR3022-IgG1] by ELISA. *Sample:* Plate coated with SARS-CoV-2 SRBD/HIS Tag (P170-100 lot 1) and Delta Variant RBD at 2 ug/ml. Antibodies were titrated on a 3-fold dilution series beginning at 5 ug/ml. *Detection:* Gt x-Human IgG Fc HRP, A80-104P at 1:50K