



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# CD79A Recombinant Monoclonal Antibody [BLR233K]

Rabbit Recombinant Monoclonal

Purified		RefSeq ID	NP_001774.1
Catalog No.	A700-233	Uniprot ID	P11911
Lot No.	A700-233-1	GeneID	12518

**APPLICATIONS** WB, IP, IHC, ICC, Flow Cyt

**SPECIES REACTIVITY** Mouse

**AMOUNT** 100 µl (50+ tests)

**CONCENTRATION** 250 µg/ml

**STORAGE/SHELF LIFE** 2 – 8°C / 1 year from date of receipt

**PHYSICAL STATE** Liquid

**BUFFER** Borate Buffered Saline (BBS) pH 8.2 with 0.1% BSA and 0.09% Sodium Azide

**ISOTYPE** IgG

**CLONE #** BLR233K

**ORIGIN** USA

**PRODUCTION PROCEDURES** Recombinant antibody was purified from cell culture supernatant.

Immunogen was a peptide representing a region between residue 176 and the C-terminus (residue 226) of human B-cell antigen receptor complex-associated protein alpha chain (aka CD79A) using the numbering given in entry NP\_001774.1 (Gene ID 12518).

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000

Immunoprecipitation 20 µl/1 mg lysate

Immunohistochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.

Immunocytochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE cell sections.


Flow Cytometry Fixed in 4% formaldehyde and permeabilized with 90% methanol. 1 µl per 1 x 10<sup>6</sup> cells.

**IHC MOUSE CONTROLS** Spleen, CH27 Cells, WEHI-231 Cells


**ADDITIONAL INFO** <https://www.fortislife.com/p/A700-233>

Use the link above to view SDS, a current list of citations, and other product specific information.


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Michael Spencer, PhD Date: September 29, 2022



**Detection of mouse CD79A by western blot.** *Samples:* Whole cell lysate (10 µg) from WEHI-231, NIH 3T3, CH27, TCMK-1, and A20 cells prepared using NETN lysis buffer. *Antibody:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR233K] (A700-233 lot 1) used at 1:1000. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 30 seconds. Lower Panel: Rabbit anti-Actin recombinant monoclonal antibody [BLR057F] (A700-057).




**Detection of mouse CD79A by immunohistochemistry.** *Sample:* FFPE section of mouse spleen. *Antibody:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR233K] (A700-233-1). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).



**Detection of mouse CD79A by western blot of immunoprecipitates.** *Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from CH27 cells prepared using NETN lysis buffer. *Antibodies:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR233K] (A700-233 lot 1) used for IP at 20 µl/mg lysate. CD79A was also immunoprecipitated by rabbit anti-CD79A antibody Brother. For blotting immunoprecipitated CD79A, A700-233 was used at 1:1000. Chemiluminescence with an exposure time of 30 seconds.



**Detection of mouse CD79A by immunocytochemistry.** *Sample:* FFPE section of WEHI-231 cells. *Antibody:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR233K] (A700-233-1). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P).



**Detection of mouse CD79A (shaded) in A20 cells by flow cytometry.** *Antibody:* Rabbit anti-CD79A recombinant monoclonal antibody [BLR233K] (A700-233) or isotype control (unshaded). *Secondary:* DyLight® 650-conjugated goat anti-rabbit IgG (A120-101D5).